



**BJP**

**Bangladesh Journal of Pharmacology**

**Research Article**

**Chemical composition and antimicrobial properties of different basil essential oils chemotypes from Togo**

## Chemical composition and antimicrobial properties of different basil essential oils chemotypes from Togo

Koffi Koba<sup>1</sup>, P.W. Poutouli<sup>2</sup>, Christine Raynaud<sup>3</sup>, Jean-Pierre Chaumont<sup>4</sup> and Komla Sanda<sup>1</sup>

<sup>1</sup>Unité de Recherche sur les Agroressources et la Santé Environnementale, Ecole Supérieure d'Agronomie, Université de Lomé, BP. 1515, Lomé Togo; <sup>2</sup>Département de Biologie Animale et de zoologie, Faculté des Sciences, Université de Lomé, B.P. 1515 Lomé Togo; <sup>3</sup>Laboratoire de Chimie Agro-Industrielle, UMR 1010, INRA/INP-ENSIACET, 118, route de Narbonne, 31077 Toulouse cedex, France; <sup>4</sup>Equipe des Sciences Séparatives et Biopharmaceutiques EA 3924, Université de Franche-Comté, Faculté de Médecine et de Pharmacie, Place Saint-Jacques, 25030, Besançon, Cedex France.

### Article Info

Received: 30 July 2008  
Accepted: 4 August 2008  
Available Online: 9 August 2008  
DOI: 10.3329/bjp.v4i1.998

### Cite this article:

Koba K, Poutouli PW, Raynaud C, Chaumont JP, Sanda K. Chemical composition and antimicrobial properties of different basil essential oils chemotypes from Togo. Bangladesh J Pharmacol. 2009; 4: 1-8.

### Abstract

The aerial parts essential oils of *Ocimum basilicum* (Lamiaceae) from Togo were steam-distilled and investigated for their percentage composition (GC and GC/MS) and *in vitro* antimicrobial activities. Five oil chemotypes were identified and classified as follows in line with their principal components: Estragole type; linalool/estragole type; methyleugenol type; methyleugenol/t-anethole type; t-anethole type. The *in vitro* microbiological experiments revealed that only the methyleugenol and methyleugenol/t-anethole chemotypes were active against tested fungi and bacteria. Their minimum inhibitory concentrations (MIC) ranged from 80-150 µL/L and from 200-500 µL/L respectively. Likewise, on tested bacteria the MIC varied from 200-400 µL/L and from 250-500 µL/L respectively. These findings are supportive of the potential of both basil oil chemotypes for use as active ingredients in natural antibiotic drugs.

### Introduction

The genus *Ocimum* collectively called basil (in English), includes around 30 plant species from tropical and subtropical areas (Paton, 1992). *Ocimum* is widely cultivated and extensively used for food, perfumery, cosmetics, pesticides, medicine, and traditional rituals because of their natural aroma and flavor and other properties (Alburquerque, 1996; Darrah, 1974). Literature reports that *O. basilicum* leaf essential oils or leaf powder have effective insecticidal and pesticidal activities against *Vigna unguiculata* pests *Callosobruchus maculatus* (F) (Coleoptera: Bruchidae) (Keita et al., 2001; Ketoh et al., 2002). *O. basilicum* commonly grows semi-wild and is cultivated in Togo at small scale in vegetable gardens. Current domestic uses are only for food and folk medicine (Adjanooun et al., 1986).

Nevertheless, a large scale production of the basil essential for well-known value-added applications (Berrada et al., 1987; Archtander, 1994) is quite viable under local circumstances.

The chemical composition of *O. basilicum* essential oils has been intensively investigated throughout the world (Ekundayo et al., 1987; Sanda et al., 1998; Yayi et al., 2001), indicating that the estragole chemotype and the linalool-estragole one are the most widely distributed.

At the same time, only very little work has been done on the chemical composition and the antimicrobial activities of basil oils from plants growing in Togo (Sanda et al., 1998; Koumaglo et al., 1996). In addition, there are very few data available either for practical use or for basic research needs about antimicrobial properties of the essential oils of *O. basilicum* growing



in Togo (Baba-Moussa et al., 1997; Chaumont et al., 2001).

In order to help fill this deficiency, partly at least, the aim of the present work was to investigate basil oils extract of plants sampled in different localities belonging to different ecological areas of Togo. Specifically, oils percentage composition was determined along with their bacteriostatic and fungistatic activity using 14 bacterial and 14 fungal strains.

The principal finding expected from this applied research is the candidate basil oil(s) that could be produced in Togo for use as source(s) of active ingredients in natural antibiotics or cosmetics.

## Materials and Methods

### *Plant material sampling and volatile oils isolation*

Leaves and inflorescences of *O. basilicum* used in this work were harvested from plants at full flowering stage in vegetable gardens in five various locations of Togo from May to October 2007.

Plant specimen was identified by Prof. Akpagana, Département de Botanique, Faculté des Sciences at the Université de Lomé (Togo), where voucher specimens were deposited in the herbarium.

A sample (50 g) of air-dried plant material was extracted by the hydrodistillation technique during 2 hours in a modified Clevenger-type glass apparatus (Craveiro et al., 1976). The extracted crude essential oils were stored in hermetically sealed dark glass flasks with rubber lids, covered with aluminium foil to protect the contents from light and kept under refrigeration at 4°C until use without any prior purification.

### *Essential oil analyses by gas chromatography*

Gas chromatographic analysis was carried out on a Varian 3300 type gas chromatograph equipped with FID detector. An apolar capillary column DB-5 (30 m x 0.25 mm i.d.; film thickness 0.25 µm) and on a polar column supelcowax 10 with the same characteristics as above mentioned were used. DB-5 column operating conditions were as follows: From 50°C (5 min), 50 to 250°C at the rate of 2°C/min and supelcowax 10 from 50°C (5 min), 50 to 200°C at 2°C/min. The injector and detector temperatures were respectively 250 and 300°C. The carrier gas was helium at a flow rate of 1.5 mL/min. Samples (0.2 µL) of undiluted essential oil were injected manually.

### *Gas chromatography-mass spectrometry analysis*

The GC/MS analysis was carried out on a Hewlett

Packard 5890 series II chromatograph, coupled with a mass spectrometer of the Hewlett Packard 5971 series type operating in the EI mode at 70 eV. The capillary column type was DB5-MS (30 m x 0.25 mm i.d.; film thickness 0.25 µm). The amount of sample injected and GC/MS parameters were the same as above.

### *Identification of components*

The components of oils samples were identified by their retention time, retention indices relative to C<sub>8</sub>-C<sub>24</sub> *n*-alkanes, computer matching with Wiley 275L library and as well as by comparison of their mass spectra with the authentic samples or with data already available in the literature (Kondjoyan and Berdagué, 1996; Adams, 2001).

The percentage of composition of the identified compounds was computed from the GC peak area without any correction factor and was calculated relatively.

### *In vitro antifungal testing*

Fungal strains used are listed in Table I. They were supplied by Institut Pasteur de Paris and Hôpital Saint Jacques de Besançon, France. The fungi were cultivated on a sabouraud agar medium in which was added chloramphenicol 1%, all purchased from BioMerieux Co. (Paris, France). Pure estragole, linalool and methyleugenol commercial standards were also purchased from BioMerieux Co. (Paris, France).

The antimicrobial activities of the essential oils were assessed according to agar dilution method (Benjilali et al., 1986; Griffin et al., 2000). The tested essential oil and its pure major components from commercial origin were diluted in a minimal quantity of ethanol 95% 1/10 v/v to which was added an aqueous solution of Tween 80 (final concentration of 1% v/v) in order to obtain a homogeneous mixture. The later was incorporated as appropriate to the microbiological culture medium under solidification to obtain final concentrations of the active ingredient that ranged from 10 to 500 µL/L. The mixture was then poured into 3 cm diameter petri dishes.

After solidification fungal strains were respectively seeded as described below:

- (i) dermatophytes were seeded with a disc of approximately 2 mm, from a mycelia carpet of pre-culture, laid in the middle area of a new petri dishes, upper side against the new culture medium;
- (ii) 1 mL of a suspension of 10<sup>5</sup> conidia per mL of *Aspergillus fumigatus* or 10<sup>5</sup> blastospores per mL of yeast was poured on the surface of the culture medium. Incubation time and temperatures depended on the fungal strains: 24 hours at 37°C for *Candida albicans* and *Aspergillus*, 48 hours at 37°C for *Cryptococcus*, 14 days at

Table I								
Antifungal activities of different <i>O. basilicum</i> essential oils chemotypes from Togo								
Fungal strains	MIC* (mL/L)							
	Basil oils chemotypes					Linalool	Méthyl eugenol	Estragole
	Ob <sup>1</sup> *	Ob <sup>2</sup> *	Ob <sup>3</sup> *	Ob <sup>4</sup> *	Ob <sup>5</sup> *			
<b>Dermatophytes</b>								
<i>Trichophyton mentagrophytes</i> (B)	>500	400	80	250	>500	>500	60	>500
<i>T. interdigitale</i> (B)*	>500	450	100	300	>500	>500	150	>500
<i>T. rubrum</i> (B)	>500	400	150	300	>500	>500	150	>500
<i>T. soudanense</i> (B)	>500	500	100	200	>500	>500	300	>500
<i>T. violaceum</i> (B)	500	500	100	300	>500	450	300	500
<i>Microsporum canis</i> (CIP)*	>500	400	200	300	>500	600	150	>500
<i>Microsporum gypseum</i> (CIP)	>500	500	200	250	500	500	150	>500
<i>Epidermophyton floccosum</i> (B)	>500	500	200	300	>500	500	200	>500
<b>Imperfect filamentous fungi</b>								
<i>Aspergillus fumigatus</i> (B)	>500	>500	150	500	>500	>500	600	>500
<i>Scopulariopsis brevicaulis</i> (B)	>500	>500	100	>500	>500	>500	500	>500
<i>Scytalidium dimidiatum</i> (B)	>500	>500	150	>500	>500	>500	>500	>500
<i>Scytalidium hyalinum</i> (B)	>500	>500	200	>500	>500	>500	>500	>500
<b>Pathogenic yeasts</b>								
<i>Candida albicans</i> (B)	>500	>500	80	500	>500	>500	100	>500
<i>Cryptococcus neoformans</i> (B)	>500	>500	100	400	>500	>500	100	>500

\*MIC: Minimum inhibition concentration; Ob<sup>1</sup>: Estragole chemotype; Ob<sup>2</sup>: Linalol/estragole chemotype ; Ob<sup>3</sup>: Methyleugenol chemotype; Ob<sup>4</sup>: Methyleugenol/t-anethole chemotype; Ob<sup>5</sup>: t-anethole chemotype; B: Hôpital Saint Jacques de Besançon, France; CIP: Collection Institut Pasteur de Paris

24°C for the dermatophytes and *Scopulariopsis brevicaulis*.

#### Antibacterial testing

Bacterial strains were supplied by Institut Pasteur Collection de Paris and Leeds University Microbiology Laboratory, United Kingdom. They were cultivated on 1.3% (m/v) nutrient broth-agar-granulated (NB-AG) culture medium (AES Laboratories), in 9 cm diameter petri dishes. After cooling and solidification, the petri dishes were automatically inoculated with the bacterial suspensions 10<sup>5</sup> CFU/mL with STEERS apparatus. The petri dishes were incubated at 37°C for 48 hours under aerobic conditions.

Major constituents: Others meaningful constituents were estragole (10.0%), (E)- $\alpha$ -bergamotene (3.6%) and linalool (2.9%).

v) t-Anethole chemotype (sample from Bassar): This sample contained mainly t-anethole (74.6%), linalool (17.3%) and carvacrol (2.7%).

To our knowledge, the last three basil oil chemotypes have not yet been reported. *O. basilicum* has been thoroughly investigated with regards to volatile oil composition but some of our findings are supportive of

the idea that continuing chemical inventory on this species is still of scientific interest to support action aimed at plant biodiversity knowledge, conservation and sustainable exploitation.

The findings on the chemical composition of basil oils in Togo were very instructive with regard to the possible applications. Hence, the estragole chemotype found here resembled that described by Ekundayo et al. (1987) and Baritau et al. (1992), which is well-known for its valuable usage in perfumery and food (Kayibou, 1992). In addition, this chemotype has been recently indicated as a possible pesticide against larva and adult of *Callosobruchus maculatus* known as a major post-harvest pest of *Vigna unguiculata* in storage (Keita et al., 2001; Ketoh et al., 2002).

The pesticidal activity of the volatile oil of *Clausena anisata*, containing mainly estragole and t-anethole (Moudachirou et al., 1997; Okunde and Olaiya, 1987). Such indications on the pesticidal properties of this essential oil containing estragole and t-anethole is certainly indicative of the biocide potential of these natural molecules which might be endowed with antimicrobial properties (Friedman et al., 2002), our focus in this investigation.

imperfect filamentous fungi were also highly sensitive to the test volatile oil; MICs were in the range of 100-150  $\mu\text{L/L}$ . Likewise, the methyleugenol oil chemotype appeared very toxic to pathogenic yeast strains including *Candida*, *Cryptococcus*, *Aspergillus*, and *Scopulariopsis*, MICs varying from 80 to 100  $\mu\text{L/L}$ . Also noticeable was the antifungal effect (MIC: 150  $\mu\text{L/L}$  and 200  $\mu\text{L/L}$ ) of this oil chemotype on both strains of *Scytalidium sp.* The later, which often resist conventional antibiotics, are usually reported in human dermatomycosis in tropical and subtropical countries (Alvarez et al., 2000). The African *Trichophyton soudanense*, a parasite frequent in school environment (Ouaffak et al., 2001; Vandemeulebroucke et al., 1999) was interestingly also sensitive to the tested essential oils (MICs 100 and 200  $\mu\text{L/L}$ ). It was also the case for *Cryptococcus neoformans*, a hazardous opportunist yeast, which is a resistant germ usually infecting humans affected by HIV/AIDS, which group of patients is known to be generally at high-risk with regard to mycosis opportune affections.

The high antifungal activity of this basil oil chemotype on those pathogenic fungi like dermatophytes, filamentous fungi and yeasts confirmed the excellent fungal growth inhibition properties previously reported as a characteristic of essential oils rich in methyleugenol and/or other phenol derivatives (Chaumont and Leger, 1989). In this work there is no doubt that the antifungal activity of *O. basilicum* chemotype methyleugenol against tested fungi is a predictable consequence of its high content in methyleugenol known as one of the phenolic volatile molecules endowed with antimicrobial properties (Chaumont and Leger, 1989; Viollon and Chaumont, 1994). In comparison, commercial linalool and estragole tested as standards in this study were found non-effective unlike pure methyleugenol standard.

The methyleugenol/t-anethole chemotype showed a moderate antifungal activity only on dermatophytes with MICs ranged from 250-300  $\mu\text{L/L}$  but not filamentous fungi and yeasts were not affected. The estragole type, the linalool/estragole type, and the t-anethole type were ineffective on all fungal strain tested in this work. A lesson learned from the findings mentioned above was that estragole and t-anethole, which are reported as possible natural pesticides, could not serve as natural fungicidal ingredients unlike methyleugenol against the fungal strains tested in this work. This simply indicates the broader understanding of what is generically termed to as the biological activity of essential oils.

Table III shows that all feet microflora bacterial strains tested were only sensitive to methyleugenol and methyleugenol/t-anethole oil chemotypes.

The MICs, which were in the range of 200-500  $\mu\text{L/L}$

were slightly higher than those recorded on fungal strains in this study, indicating a lesser activity. The results recorded against common armpit microflora bacterial strains, were especially promising for a possible use against *Corynebacterium xerosis*, the principal one of the bacteria responsible for bad odors. Its growth was interestingly inhibited with at a dose of 100  $\mu\text{L/L}$  antibacterial test suspension. Note that this bacteria metabolizes human steroids produced via transpiration to yield small and bad-smelling volatile molecules (Rennie et al., 1991).

The findings in this study indicated bacteriostatic effects against feet microflora germs like *Staphylococcus epidermidis* and *Staphylococcus hominis* (MIC: 200 and 300  $\mu\text{L/L}$ ) partly responsible for feet bad odors (Marshall et al., 1987; Marshall et al., 1988). Methyleugenol and methyleugenol/t-anethole oil chemotypes samples were also very active against the *Coryneform* (*Corynebacterium* gr B, C and D2) (MIC: From 100-300  $\mu\text{L/L}$ ) which produce proteinase and lipase, both enzymes that cause feet surface skin bad odors (Marshall et al., 1987; Marshall et al., 1988). Both oil chemotypes also showed a moderate antibacterial activity against gram negative bacteria (*Acinetobacter sp*, *Moraxella sp*: MIC from 250- 500  $\mu\text{L/L}$ ) and gram positive bacteria (*Micrococcus luteus*, *Staphylococcus epidermidis*: MIC: From 200-500  $\mu\text{L/L}$ ). The later are involved in nosocomial infections (Bergogne-Bérézín, 1995; CDC, 1999).

The proven antibacterial activities of the methyleugenol and metyleugenol/t-anethole types of basil oil found in this study are quite indicative of their potential as possible active ingredients for use in the formulation of deodorants for armpits or feet. Developing sustainable large scale consumer goods like natural drugs and cosmetics based on basil oil as well as on other essential oils bearing aromatic plants growing in Togo is of an obvious economic interest while helping preserve plant biodiversity.

Culture media with or without ethanol 95% and Tween 80 were used as controls.

The antimicrobial activities were evaluated by the determination of the minimum inhibitory concentration (MIC). The MIC of a tested active ingredient was determined as the lowest concentration of the test antimicrobial ingredient sample that resulted in a complete inhibition of visible growth of the microorganisms. All tests were carried out in triplicate.

---

## Results and Discussion

Basil oil extraction yields were in the range of 1.4% to 2.2%, based on plant material dry weight. Identified oil constituents and percentage and their relative

Table II						
Chemical composition of <i>O. basilicum</i> volatile oils from Togo						
Identified compounds*	Peak area [%] of plant material sampling localities					
	RI <sup>a</sup>	Lomé	Lo-J.B <sup>b</sup>	Adéticopé	Bassar	Sokodé
Monoterpene hydrocarbons		3.0	12.9	1.7	0.8	4.2
α-Pinene	941	0.2	0.5			0.4
Sabinene	976	0.3	0.6			
Myrcene	993	0.3	0.6			0.4
P-Cymene	1030		0.9			0.5
Limonene	1033	0.2	6.2	1.5	0.8	0.4
(Z)-β-Ocimene	1046		3.2			0.7
(E)-Cis-ocimene	1058	1.8		0.2		
γ-Terpinene	1068	0.2				0.2
Terpinolene	1100	0.2	1.0			1.5
Oxygenated monoterpenes		6.0	6.0	3.4	22.2	49.1
1,8 Cineole	1023	2.3				3.6
Linalool	1099	1.7		2.9	17.3	41.2
Camphor	1146	0.7		0.5	1.3	0.5
Terpineol-4	1171	0.5				2.5
Estragole	1198	85.5		10.0		22.2
t-Anethole	1253			32.6	74.6	
Carvacrol	1273				2.7	
Thymol	1290				0.5	
Bornyl acetate	1289	0.2			0.5	0.9
Methyleugenol	1293	0.4	74.5	42.3		
Geranyl formiate	1381	0.5				0.4
Sesquiterpene hydrocarbons		3.7	5.3	6.3	2.3	13.5
β-Elemene	1387	0.4		0.3		1.2
β-Caryophyllene	1420	0.4				0.2
(E)-α-Bergamotene	1440	1.6		3.6	0.6	7.6
α-Caryophyllene	1452			0.3		0.4
Germacrene D	1487	0.3	0.6	0.2		0.6
β-Selinene	1493			0.2		0.8
α-Zingiberene	1494			0.4		
Bicyclogermacrene	1502	0.1				0.4
α-Muurolene	1508	0.3	0.6			0.9
Germacrene A	1513	0.6		0.4		0.1
γ-Cadinene	1514	0	4.1	0.9	1.7	1.4
Oxygenated sesquiterpenes		0.5		3.7		0.4
Γ-Cadinol	1619	0.5		3.7		0.4
Total identified (%)		98.6	92.3	99.7	100.0	89.4

The experimental data in Table II show that all fungal strains tested were sensitive to the metyleugenol type of basil oil, but dermatophytes and pathogenic yeasts

were more particularly affected. Hence, markedly low MICs (80 μL/L) were recorded with *T. mentagrophytes var mentagrophytes*. Along with dermatophytes,

Table III

## Antibacterial activities of different basil essential oils chemotypes from Togo

Cutaneous bacterial strains	MIC <sup>a</sup> (mL/L)							
	Basil oils chemotypes					Linalol	Méthyl eugénol	Estragole
	Ob <sup>1b</sup>	Ob <sup>2</sup>	Ob <sup>3</sup>	Ob <sup>4</sup>	Ob <sup>5</sup>			
Feet microflora								
<i>Staphylococcus epidermidis</i> (L1S2) <sup>c</sup>	>500	>500	200	300	>500	>500	>500	>500
<i>S. hominis</i> (L8S2) <sup>c</sup>	>500	>500	200	400	>500	>500	>500	>500
<i>S. cohnii</i> * (L6S3)	>500	>500	150	300	>500	>500	>500	>500
<i>Coryneform</i> gr. B (L 16C3) <sup>c</sup>	>500	400	100	300	>500	>500	>500	>500
<i>Coryneform</i> gr. C (L3C3) <sup>c</sup>	>500	>500	150	250	>500	>500	>500	>500
<i>Coryneform</i> gr. D2 (L19C1) <sup>c</sup>	>500	500	150	200	>500	>500	>500	>500
<i>Micrococcus sedentarius</i> (L7B5) <sup>c</sup>	>500	400	200	300	500	>500	>500	>500
<i>Acinetobacter</i> sp. (LLH5DC1)	>500	500	300	400	400	>500	500	>500
<i>Moraxella</i> sp. (LH7SV1)	>500	500	250	500	>500	400	>500	>500
<i>Alcaligenes</i> sp. (LH4TV1)	>500	400	300	300	500	>500	>500	>500
Armpits microflora								
<i>Staphylococcus xylosum</i> (IP 8166)	>500	>500	300	400	>500	>500	>500	>500
<i>S. haemolyticus</i> (IP 8156) <sup>c</sup>	>500	>500	200	300	>500	>500	>500	>500
<i>Corynebacterium xerosis</i> (IP5216) <sup>c</sup>	>500	>500	150	250	>500	>500	>500	>500
<i>Micrococcus luteus</i> (L1C5) <sup>a</sup>	>500	>500	400	500	>500	>500	>500	>500

<sup>a</sup>MIC: Minimum inhibitory concentration; <sup>b</sup>Ob<sup>1</sup>: Estragole chemotype; Ob<sup>2</sup>: Linalol/estragole chemotype; Ob<sup>3</sup>: Methyleugénol chimiotype; Ob<sup>4</sup>: Methyleugenol/t-anethole chemotype; Ob<sup>5</sup>: t-anethole chemotype<sup>c</sup>: Bacteria strains responsible of bad odours; IP: Bacteria strains provided by Institut Pasteur de Paris; L, LH: Bacteria strains provided by Dr. Marshall, Leeds University Microbiology Laboratory (United Kingdom)

percentages are listed in Table II.

Using the relative importance of the oils major constituents as classification criteria permitted to determine five chemotypes as listed below:

i) Estragole chemotype (sample from Lomé): It contained mainly estragole (85.50%), with a little amount of 1,8-cineole (2.25%). The estragole content in this sample was similar to previous findings (Guenther, 1949; Baritoux et al., 1992; Yayi et al., 2001).

ii) Linalol-estragole chemotype (sample from Sokodé): this type of basil oil contained linalool (41.21%), estragole (22.17%) and (E)- $\alpha$ -bergamotene (7.56%) as major constituents. This chemotype was very close to the European type previously reported by other workers in Nigeria (Ekundayo et al., 1987) and in Benin (Yayi et al., 2001).

iii) Methyleugenol chemotype (sample from Botanical garden, UL): This basil oil type contained mainly methyleugenol (74.45%), limonene (6.22%),  $\gamma$ -cadinene (4.07%) and (Z)- $\beta$ -ocimene (3.16%).

Antibacterial activities are shown in Table III.

## Conclusion

The prospective investigation of the percentage composition of *O. basilicum* essential oils along with the evaluation of their antimicrobial properties on human normal cutaneous microflora bacterial and fungal strains is quite a typical applied research. The ultimate goal of this type of investigation is to help protect plant biodiversity and validate on a scientific basis the potential of some valuable species whether for food or non-food applications. The present study has shown that it is feasible to use some kinds of the leaf essential oil of *O. basilicum* growing in Togo as a natural powerful antimicrobial ingredient whether in traditional or in modern medicines as well as in cosmetics. Such oils are the methyleugenol and the methyleugenol/t-anethole types. Conversely, the estragole type, the linalool/estragole type, and the t-anethole type, which were also included in the investigated samples, did show any useful antimicrobial properties.

## Financial Support

Self-funded

## Conflict of Interest

Authors declare no conflict of interest

## References

- Adams RP. Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy. Carol Stream, Illinois, Allured Publishing Corporation, 2001.
- Adjanohoun E, Ahyi MRA, Aké Assi L, Akpagana K, Chibon P, El-Hadj A, Eymen I, Goutote E, Ginko S, Hodouto KK, Hougnon P, Keita A, Kéoula Y, Klouga-Ocloo WP, Lo I, Siamevi K, Taffame KK, Garba M, Gassita JN, Gbeassor M. Médecine traditionnelle et Pharmacopée: Contribution aux études ethno-botaniques et floristiques du Togo. ACCT, Paris, 1986.
- Albuquerque UP. de Taxonomia e etnobotânica do gênero *Ocimum* L. (Lamiaceae) no Nordeste do Brasil-renfêrencia especial para Pernambuco. Recife. Dissertação (Mestrado em Biologia Vegetal)- Centro de Ciencias Biológicas, Universidade Federal de Pernambuco, 1996, p 125.
- Archander S. Perfume and flavour material of natural origin. Carol Steam, Allured Publishing Corporation, 1994, p 81.
- Alvarez P, Enriquez AM, Toro C, Martinez I, Buhigas I, de Miguel S, Lago M, Puente S, del Palacio A, Baquero M. Dermatomiucosis de importación por *Scytalidium dimidiatum*: A propósito de tres casos. Rev Iberoam Micol. 2000; 17: 102-06.
- Baba-Moussa F, Koumaglo K, Ayédoun K, Akpagana K, Moudachirou M, Bouchet P. Activité antifongique d'huiles essentielles extraites au Bénin et au Togo. Cryptogamie Mycol. 1997; 18: 165-68.
- Baritau O, Richard H, Touche J, Derbesy M. Effects of drying and storage of herbs and spices on the essential oil. Part I. Basil, *O. basilicum* L. Flav Fragr J. 1992; 7: 267-71.
- Benjilali B, Tantaoui-Elaraki A, Ismaïlou-Alaoui M, Ayadi A. Méthode d'étude des propriétés antiseptiques des huiles essentielles par contact direct en milieu gélosé. Plant Méd Phytother. 1986; 20: 155-67.
- Berrada M, Ait Igrri M, Il Idrissi A, Bellakhdar. Contribution à l'analyse de l'huile essentielle d'un cultivar d'*Ocimum basilicum* L. du Maroc. Al Buruniya. Rev Mar Pharm. 1987; 3: 137-44.
- Bergogne-Bérézin E. The increasing significance of outbreaks of *Acinetobacter* spp: The need for control and new agents. J Hosp Infect. 1995; 30: 441-52.
- Chaumont JP, Leger D. Propriétés antifongiques de quelques phénols et de composés chimiquement très voisins. Relations structure-activité. Plant Med Phytother. 1989; 23: 124-28.
- Chaumont JP, Mandin D, Sanda K, Koba K, De Souza C. Activités antimicrobiennes de cinq huiles essentielles de lamiacées togolaises vis à vis de germes représentatifs de la microflore cutanée. Acta Bot Gallica. 2001; 148: 93-101.
- Center for Disease Control and Prevention (CDC). National nosocomial infections surveillance system report. Data summary from January 1990-May 1990. Am J Infect Control. 1999; 27: 520-32.
- Craveiro AA, Matos FJ, Alencar JW. A simple and inexpensive steam generator for essential oils extraction. J Chem Educ. 1976; 53: 652.
- Darrah HH. Investigation on cultivars of Basils (*Ocimum*). Econ Bota. 1974; 28: 63-67.
- Ekundayo O, Laakso I, Oguntimein B, Okogun JL, Elujoba A, Hulbinen A. Essential oil of *Ocimum basilicum* from Nigeria. Acta Pharm Fen. 1987; 96: 101-06.
- Friedman M, Henika PR, Mandrell RE. Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella enterica*. J Food Prot. 2002; 65: 1545-60.
- Griffin SG, Markham JL, Leach DN. An agar dilution method for the determination of the minimum inhibitory concentration of essential oils. J Essent Oil Res. 2000; 12: 249-55.
- Guenther E. The essential oils. Vol. 2. New York, Van Nostrand, 1949, pp 16-18, 503-05.
- Kayibou G. Les Eucalyptus du Congo. Variations inter et intra-spécifiques du rendements et de la composition de leurs huiles essentielles. Thèse de doctorat Université de Montpellier II, France, 1992, p 112.
- Ketoh GK, Glitho AI, Huignard J. Susceptibility of the bruchus *Callosobruchus maculatus* (Coleoptera: Bruchidae) and its parasitoid *Dinarmus basalis* (Hymenoptera: Pteromatidae) to three essential oils. J Econ Entomol. 2002; 95: 174-82.
- Keita SM, Vincent C, Schmit J, Arnason JT, Belanger A. Efficacy of oil of *Ocimum basilicum* L. and *O. gratissimum* L. applied as an insecticidal fumigant and powder to control *Callosobruchus maculatus* (Fab). J Stored Prod Res. 2001; 37: 339-49.
- Kondjoyan N, Berdagué JL. A compilation of relative retention indices for the analysis of aromatic compounds. Ed. Laboratoire Flaveur, INRA de Theix, France, 1996.
- Koumaglo K, Dotse K, Akpagana K, Garneau FX, Gagnon H, Jean IF, Moudachirou M, Addae-Mensah I. Analyse des huiles essentielles de deux plantes aromatiques du Togo. Riv Ital EPPOS. 1996; 7: 680-91.
- Marshall J, Leeming JP, Holland KT. The cutaneous microbiology of normal human feet. J Appl Bacteriol. 1987; 62: 139-46.
- Marshall J, Holland KT, Gribbon EM. A comparative study of cutaneous microflora of normal feet with low and high levels of odour. J Appl Bacteriol. 1988; 65: 61-68.
- Moudachirou M, Ayedou MA, Gbenou J, Garneau FX, Koumaglo KH, Mensah-Addae I. Composition chimique des huiles essentielles des feuilles de *Clausena anisata* récoltées dans la sous-region Bénin-Togo-Ghana. J Soachim. 1997; 3: 49-54.
- Okunde AL, Olaifa JL. Estragole: An acute toxic principle from the volatile oil of the leaves of *Clausena anisata*. J Nat Prod. 1987; 50: 990-91.



- Ouaffak L, Gati A, Lyagoubi M. Les teignes du cuir chevelu dans les écoles Primaires de Khermisset (Maroc). *J Mycol Méd.* 2001; 11: 181-84.
- Paton A. A synopsis of *Ocimum* L. (Labiatae) in Africa. *Kew Bull.* 1992; 47: 403-35.
- Rennie PJ, Gower DB, Holland KT. *In vitro* studies of human axillary odour and cutaneous microflora. *Br J Dermatol.* 1991; 124: 596-602.
- Sanda K, Koba K, Nambo P, Gaset A. Chemical investigation of *Ocimum* species growing in Togo. *Flav Fragr J.* 1998; 13: 226-32.
- Vandemeulebroucke E, Mounkassa B, Loye J, Jousserand P, Poujade F, Petithory JC. Teignes du cuir chevelu en milieu scolaire rural au Mali. *J Mycol Med.* 1999; 92: 11-13.
- Viollon C, Chaumont JP. Antifungal properties of essential oils and their main components upon *Cryptococcus neoformans*. *Mycopathologia* 1994; 128: 151-53.
- Yayi E, Moudachirou M, Chalchat JC. Chemotyping of three *Ocimum* species from Benin: *O. basilicum*, *O. canum* and *O. gratissimum*. *J Essent Oil Res.* 2001; 13: 13-17.
- 

**Author Info**

Koffi Koba (Principal contact)  
e-mail: danielkkoba@yahoo.fr