

BJP

Bangladesh Journal of Pharmacology

Research Article

***In vitro* glucose uptake by isolated
rat hemi-diaphragm study of *Aegle
marmelos* Correa root**

In vitro glucose uptake by isolated rat hemi-diaphragm study of *Aegle marmelos* Correa root

Subban Ravi¹, C. T. Sadashiva², T. Tamizmani³, T. Balasubramanian³, M. Rupeshkumar³ and Indira Balachandran²

¹Department of Chemistry, Karpagam University, Coimbatore, Tamilnadu, India; ²Phytochemistry Division, Centre for Medicinal Plants Research, Arya Vaidya Sala, Kottakkal 676 503, Kerala, India; ³Bharathi college of Pharmacy,

Article Info

Received: 15 October 2008

Accepted: 2 November 2008

Available Online: 5 November 2008

DOI: 10.3329/bjp.v4i1.1211

Cite this article:

Ravi S, Sadashiva CT, Tamizmani T, Balasubramanian T, Rupeshkumar M, Balachandran I. *In vitro* glucose uptake by isolated rat hemi-diaphragm study of *Aegle marmelos* Correa root. Bangladesh J Pharmacol. 2009; 4: 65-68.

Abstract

The methanol extract of the root of *Aegle marmelos*, a medicinal plant, was fractionated into eight fractions using column chromatography. The antidiabetic activity of all the fractions was studied using the glucose uptake by isolated rat hemi-diaphragm *in vitro* model. Using the bioassay-guided fractionation, two compounds **1** and **2** were isolated by column chromatography and identified as 6-methyl-4-chromanone and skimmianine respectively by NMR and mass spectral methods.

Introduction

An increasing number of research papers and reviews clearly indicate that medicinal plants exhibit a variety of therapeutic properties and could provide health security to rural people primary health care. Among medicinal plants *Aegle marmelos* Correa (family Rutaceae) appears to be relevant and this plant is available in India, Bangladesh, Burma and Sri Lanka. Its distribution is mainly within the sub-Himalayan forests, in dry hilly places ascending to 4,000 feet. It is called "Shivadume", the tree of Shiva. Since ancient time, its leaves are offered to Shiva and Parvathi. *A. marmelos* has an important place in indigenous systems of medicine. With respect to pharmacology, alcoholic and aqueous extracts of the leaves had similar effects as digoxin in amplitude and contractions of the frog heart and methanolic extracts of roots inhibited the beating rate by approximately 50% of cultured mouse myocardial cells (Kakiuchi et al., 1991). Alcoholic extracts of the roots and fruits showed hypoglycemic and antidiabetic

activity (Kamalakkannan et al., 2003; Karuna-nayake et al., 1984; Sabu and Kuttan, 2004). With respect to clinical applications, it should be noted that the roots are astringent, bitter and febrifuge. They are useful in diarrhea, dysentery, dyspepsia, stomachalgia (Shoba and Thomas, 2001), cardiopalmus, seminal weakness, vomiting, intermittent fever swellings. The leaves of *A. marmelos* are useful as laxative, febrifuge and expectorant, also in ophthalmia, deafness, inflammations, cataract, diabetes, asthmatic and antifungal complaints (Rana et al., 1997). Also, the effect of these extracts was examined in the regulation of hyperthyroidism (Kar et al., 2002) and for the analgesic activity in mice. The stem extract inhibit *in vitro* proliferation of human tumor cell lines (Lampronti et al., 2003).

Diabetes mellitus is a major disease characterized by derangement in carbohydrate, fat and protein metabolism, affecting nearly 10% of the population. In the recent past many hypoglycemic agents are introduced, still the diabetes and the related complications continue



to be a major medical problem not only in developed countries but also in developing countries. Many Indian medicinal plants are reported to be useful in diabetes. However, search for new antidiabetic drugs continue. Although the leaves and fruits of *A. marmelos* were studied widely for their antidiabetic activity (Narendrakannan et al., 2006; Kesari et al., 2006), the reports on antidiabetic activity of roots are limited. The present study was aimed at investigating the antidiabetic effect of *A. marmelos* and identifying the active principles.

Materials and Methods

Plant material

A. marmelos was collected from Kottakkal, Mallapuram district, Kerala during September 2007 and authenticated at the Botany Division, Centre of Medicinal Plants Research, Arya Vaidya Sala, Kottakkal, Kerala, where the voucher specimen is maintained.

Extraction

About 20 g of air-dried root of the *A. marmelos* was powdered and refluxed with methanol for 8 hours over a water bath. The extract was filtered and concentrated under reduced pressure in a rotary evaporator below 50°C. The concentrated extract was dissolved in 10 mL of methanol and used for the gas chromatography analysis.

Fractionation using column chromatography

The air-dried root (500 g) was ground and extracted exhaustively with methanol (3 x 1.0 L) for 48 hours. The extracts were combined and freed of solvent to give 2.7 g of residue. The residue was subjected to column chromatography on 80 g of Si gel and eluted with *n*-hexane (100 mL) followed with a mixture of *n*-hexane and ethyl acetate containing increasing amounts of ethyl acetate. Finally the column was eluted with pure ethyl acetate. Seventy-eight fractions (25 mL each) were collected and pooled according to their similarity on analytical TLC plates and dried. Combined fractions 1-9 (FA, 165 mg), 10-21 (FB, 190 mg), 22-27 (FC, 141 mg), 28-35 (FD, 132 mg), 36-47 (FE, 141 mg), 48-60 (FF, 181 mg), 61-72 (FG, 156 mg) and 73-78 (FH, 153 mg) were obtained.

Isolation of compounds

The fraction FC (100 mg) was subjected to column chromatography on 20 g of Si gel G and eluted with a mixture of *n*-hexane: Ethyl acetate (8:2). Fractions (5 mL each) were collected and pooled according to their similarity on analytical TLC plates and dried. Combined fractions 7-10 (**1**, 21 mg), 18-22 (**2**, 10 mg) were obtained. They were characterized by NMR and MS

spectral data as 6-methyl-4-chromanone and skimmianine respectively.

¹H NMR (500 MHz) and ¹³C NMR (150.9 MHz) spectra were measured on a JEOL JNM LA-500 instrument using acetone-d₆ and CD₃OD solvents. APCI-MS were determined on an Agilent (Palo Alto, CA, USA) MSD 1100 single-quadrupole spectrometer.

Compound 1

APCI- MS (negative mode) m/z 161 [M-H]⁻; NMR: δ (CDCl₃) 2.35 (3H, s, methyl protons), 3.01 (2H, d, CH₂CO), 4.21 (2H, d, OCH₂), 6.78 (1H, d, aromatic) and 7.32 (2H, m, aromatic). ¹³C-NMR data; δ 66.9 (C-2), 37.8 (C-3), 196.8 (C-4), 129.2 (C-5), 128.7 (C-6), 134.10 (C-7), 114.2 (C-8), 154.2 (C-9) and 123.5 (C-10).

Compound 2

APCI-MS (positive mode) m/z 260 [M+H]⁺; NMR: δ (CDCl₃) 3.68 (9H, s, OCH₃), and aromatic protons at δ 6.4 (1H, d), 7.35(3H, m). ¹³C-NMR data; δ 149.8 (C-3), 141.8 (C-4), 118.3 (C-5), 116.9 (C-6), 154.1 (C-7), 104.2 (C-8), 146.2 (C-9) and 163.5 (C-10), 103.4 (C-11), 157.4(C-12) and 140.6 (C-13).

Gas chromatography analysis

An Agilent 6890N model GC gas chromatograph was used for the analysis. The analysis was performed as described elsewhere (Lampronti et al., 2003).

Animals

Colony bred, healthy Wister albino rats of either sex weighing 200-220 g were taken for the study. The animals were fed on standard laboratory diet with water *ad libitum* and housed at room temperature. The rats were kept of fasting overnight with free access to water during the experiment in the same ambience. The animals were divided into four groups of six animals each.

Glucose uptake by isolated rat hemi-diaphragm

Glucose uptake by rat hemi-diaphragm was estimated by the methods described elsewhere (Walaas and Walaas, 1952; Chattopadhyay et al., 1992) with some modifications. Four sets containing six numbers of graduated test tubes (n = 6) for each fraction (Fraction 1-8) were taken. Group I served as a control which contained 2 mL of Tyrode solution with 2% glucose, Group II contained 2 mL Tyrode solution with 2% glucose and regular insulin (Nova Nardisk) 0.6 mL of 0.4 units per mL solution. Group III contained 2 mL Tyrode solution with 2% glucose and 1.4 mL of *A. marmelos* fraction and the Group IV contained 2 mL Tyrode solution with 2% glucose and regular insulin 0.6 mL of 0.4 units per mL solution and 1.4 mL of *A. marmelos* fraction.

The volumes of all the test tubes were made up to 4 mL

Table I		
Effect of <i>A. marmelos</i> fractions FA-FH using glucose uptake by isolated rat-hemi diaphragm <i>in vitro</i> assay		
SN.	Group	Glucose uptake (mg/g/30 min)
1	Control	5.7 ± 0.01
2	Insulin	15.7 ± 0.2 ^b
3	FA	20.3 ± 0.2
4	Insulin + FA	19.7 ± 0.2
5	FB	30.3 ± 0.2 ^c
6	Insulin + FB	28.8 ± 0.3
7	FC	40.7 ± 0.2 ^c
8	Insulin + FC	30.7 ± 0.2
9	FD	6.8 ± 0.3
10	Insulin + FD	8.3 ± 0.2
11	FE	24.3 ± 0.2 ^c
12	Insulin + FE	22.7 ± 0.2
13	FF	22.3 ± 0.2 ^c
14	Insulin + FF	20.3 ± 0.2
15	FG	24.3 ± 0.2 ^c
16	Insulin + FG	23.2 ± 0.2
17	FH	21.2 ± 0.3 ^c
18	Insulin FH	20.2 ± 0.2 ^a

Values are mean ± SEM; n = 6; ^ap<0.05, ^bp<0.01, ^cp<0.001 as compared to the control group, one-way ANOVA

with distilled water to match the volume of the test tubes of group IV. Albino rats were tested overnight and killed by decapitation. The diaphragms were dissected out quickly with minimal trauma and divided into two halves. Two diaphragms from the same animal were not used for the same set of experiment. Six numbers of diaphragms were used for each group (Fr 1-8). The hemi-diaphragms were placed in test tubes and incubated for 30 min at 37°C in an atmosphere of 100% oxygen with shaking at 140 cycles/min. Glucose uptake per gram of tissue was calculated as the difference between the initial and final glucose content in the incubated medium.

Statistics

The data were analyzed using one-way ANOVA followed by Dunnett's test.

Results and Discussion

The estimation of glucose content in rat hemi-diaphragm was employed for *in vitro* study of peripheral uptake of glucose. The effects of the *A. marmelos* fractions FA-FH on glucose uptake by isolated rat-hemi diaphragm are shown in Table I. The fractions FB and FC enhances the uptake of glucose by isolated rat hemi-diaphragm significantly (p<0.001) and was

found to be more effective than insulin. Administration of *A. marmelos* fractions and insulin together was found to be less effective than *A. marmelos* fractions alone, but significantly higher than insulin treated group (p<0.001).

The gas chromatography analysis was performed as described in Lampronti et al. (2003). The resulting chromatogram indicated that one intense peak (retention time 19.3 min), corresponding to 6-methyl-4-chromanone, is present in the extract which matches with the results of Lampronti et al. (2003).

Fractions FB and FC were analyzed with TLC and found that there were two spots in the fractions FB and FC showing the same R_f value at 0.4 and 0.5. These two compounds 1 (13 mg) and 2 (9 mg) were isolated by column chromatography of the fraction FB on Si gel G and eluted with a mixture of *n*-hexane: Ethyl acetate (8:2). The structure of the compounds (Figure 1) was elucidated using MS and ¹H-NMR analysis.

Conventional antidiabetic agents can affect several pathways of glucose metabolism such as insulin secretion, glucose uptake by target organs as well as nutrient absorption.

The estimation of glucose content in rat hemi-diaphragm is a commonly employed and reliable method for *in vitro* study of peripheral uptake of glucose.

Among the fractions tested the fractions FB and FC enhances the uptake of glucose by isolated rat hemi-diaphragm significantly (p<0.001) and was found to be more effective than insulin. It appears that fractions FB and FC have direct peripheral action. Administration of *A. marmelos* fractions and insulin together was found to be less effective than *A. marmelos* fractions alone, but significantly higher than insulin treated group (p<0.001). It appears that drug interaction could have occurred between *A. marmelos* fractions and insulin when given together. The control value of the glucose uptake by rat hemi-diaphragm (5.7 ± 0.01) corresponds with the findings of Chattopadhyay et al. (1992). Gas chromatography analysis indicated the presence of 6-methyl-4-chromanone, in the extract which matches with the results of Lampronti et al. (2003).

Compound 1 had a molecular formulae C₁₀ H₁₀ O₂




Figure 1: Structures of compounds 1 and 2

evident from its APCI- MS (negative mode) spectrum by exhibiting a peak at m/z 161 [M-H]⁻. Its NMR spectrum showed peaks at δ 2.35 (3H, s, methyl group attached to an aromatic ring), 3.01 (2H, d, -CH₂CO), 4.21 (2H, d, -OCH₂), 6.78 (1H,d, aromatic) and 7.32 (2H, m, aromatic) and identified as 6-methyl-4-chromanone. ¹³C-NMR data confirms the structure of **1**; δ 66.9 (C-2), 37.8 (C-3), 196.8 (C-4), 129.2 (C-5), 128.7 (C-6), 134.10 (C-7), 114.2 (C-8), 154.2 (C-9) and 123.5 (C-10). Compound **2** had a molecular formulae C₁₄ H₁₃ O₄ N evident from its APCI-MS (positive mode) spectrum by exhibiting a peak at m/z 260 [M+H]⁺. Its NMR spectrum showed peaks at δ 3.68 (9H, s, three - OCH₃), and aromatic protons at δ 6.4 (1H, d), 7.35 (3H, m) and the ¹³C-NMR data confirms the structure of **2** as skimmianine; δ 149.8 (C-3), 141.8 (C-4), 118.3 (C-5), 116.9 (C-6), 154.1 (C-7), 104.2 (C-8), 146.2 (C-9) and 163.5 (C-10), 103.4 (C-11), 157.4(C-12) and 140.6 (C-13). Already skimmianine, was found to have a significant inhibitory effect on spontaneous motor activity, exploratory behavior, cataleptogenic activity, conditioned avoidance response and long-term isolation-induced fighting of animals. A mild antimethamphetamine activity was also observed. Its neuroleptic activity was less potent than that of chlorpromazine (Cheng, 2003), it has inhibitory activity against the enzyme adenine phosphoribosyltransferase (APRT) from Leishmania, a tropical parasite causing endemic disease in poor countries (Napolitano et al., 2008), function as antagonists of 5-hydroxytryptamine receptors in animals (Cheng et al., 1994), and 6-methyl-4-chromanone was earlier isolated from stem extract of *A. marmelos* and found to inhibit the *in vitro* proliferation of human tumor cell lines, including the leukemic K562, T-lymphoid Jurkat, B-lymphoid Raji, erythroleukemic HEL, melanoma Colo38, and breast cancer MCF7 and MDA-MB-231 cell lines (Lampronti et al., 2003). Our results suggest that 6-methyl-4-chromanone **1** and skimmianine **2** may be responsible for the *in vitro* antidiabetic effect.

References

- Chattopadhyay RR, Sarkar SK, Ganguly S, Banerjee RN, Basu TK. Effect of leaves of *Vinca rosea* Linn, on glucose utilization and glycogen deposition by isolated rat hemidiaphragm. *Indian J Physiol Pharmacol*. 1992; 36: 137-38.
- Cheng JT. Effect of skimmianine on animal behavior. *Acta Cryst*. 2003; E59: 1503-05.
- Cheng JT, Chang TK, Chen IS. Skimmianine and related furoquinolines function as antagonists of 5-hydroxy-tryptamine receptors in animals. *J Auton Pharmacol*. 1994; 14: 365-74.
- Kakiuchi N, Senaratne LR, Huang SL, Yang XW, Hattori M, Pilapitiya U, Namba T. Effects of constituents of *Beli (Aegle marmelos)* on spontaneous beating and calcium paradox of myocardial cells. *Plants Med*. 1991; 57: 43-46.
- Kamalakkannan N, Rajadurai M, Prince PS. Effect of *Aegle marmelos* fruits on normal and streptozotocin-diabetic Wistar rats. *J Med Food*. 2003; 6: 93-98.
- Kar A, Panda S, Barthi S. Relative efficacy of these medicinal plant extracts in the alteration of thyroid hormone concentration in male mice. *J Ethnopharmacol*. 2002; 81: 281-85.
- Karunanayake EH, Welihinda J, Sirimanne SR, Sinnadorai G. Oral hypoglycemic activity of some medicinal plants of Sri Lanka. *J Ethnopharmacol*. 1984; 11: 223-31.
- Kesari A, Gupta RK, Singh SK, Diwakar S, Watal G. Hypoglycemic and antihyperglycemic activity of *A. marmelos* seed extract in normal and diabetic rats. *J Ethnopharmacol*. 2006; 107: 374-79.
- Lampronti I, Martello N, Bianchi N, Borgatti M, Lambertini E, Piva R, Jabbar S, Choudhuri MSK, Khan MTH, Gambari R. *In vitro* anti-proliferative effects on human tumor cell lines of extracts from the Bangladeshi medicinal plant *Aegle marmelos* Correa. *Phytomedicine* 2003; 10: 300-08.
- Napolitano HB, Silva M, Ellena J, Rocha WC, Vieira PC, Thiemann OH, Oliva G. Redetermination of skimmianine: A new inhibitor against the *Leishmania* APRT enzyme. *Automatic Autacoid Pharmacol*. 2008; 14: 365-74.
- Narendrakannan RT, Subramanian S, Kandaswamy M. Biochemical evaluation of anti-diabetogenic properties of some commonly used Indian plants on streptozotocin-induced diabetes in experimental rats. *Clin Exp Pharmacol Physiol*. 2006; 33: 1150-57.
- Rana BK, Singh UP, Taneja V. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *J Ethnopharmacol*. 1997; 57: 29-34.
- Sabu MC, Kuttan R. Antidiabetic activity of *Aegle marmelos* and its relationship with its antioxidant properties. *Indian J Physiol Pharmacol*. 2004; 48: 81-88.
- Shoba FG, Thomas M. Study of anti-diarrheal activity of four plants in castor oil-induced diarrhea. *J Ethnopharmacol*. 2001; 76: 73-76.

Author Info

Subban Ravi (Principal contact)
e-mail: ravisubban@rediffmail.com