



BJP

Bangladesh Journal of Pharmacology

Research Article

Trichoderma harzianum: A new fungal source for the production of cyclosporin A

Trichoderma harzianum: A new fungal source for the production of cyclosporin A

Asma Azam, Tehmina Anjum and Wajiha Irum

Institute of Plant Pathology, University of the Punjab, Quaid-e-Azam Campus, Lahore, Pakistan.

Article Info

Received: 12 March 2012
Accepted: 28 March 2012
Available Online: 10 April 2012
DOI: 10.3329/bjp.v7i1.10084

Cite this article:

Azam A, Anjum T, Irum W. *Trichoderma harzianum*: A new fungal source for the production of cyclosporin A. Bangladesh J Pharmacol. 2012; 7: 33-35.

Abstract

Pure cultures of *Trichoderma harzianum*, *T. aureoviride*, *T. reesei*, *T. koningii* and *T. hemantum* were checked for their potential to produce cyclosporin A. Production medium used for drug production was consisted of: glucose, 5%; peptone, 1%; KH_2PO_4 , 0.5%; KCL, 0.25% (w/v). Whereas, butyl acetate was used to extract the fermentation medium for cyclosporine A. This was then analyzed through high performance liquid chromatography and the chromatograms obtained were compared with that of cyclosporine and with the external standard cyclosporin A 98.5% pure. Only chromatogram of *T. harzianum* showed a peak at 2.78, which was comparable with both the standards. Mass spectroscopy of this peak showed $[\text{CsA} + \text{H}]^+$ ion of m/z 1203. The amount of drug calculated was 44.06 $\mu\text{g}/\text{mL}$.

Introduction

Cyclosporin A is a member of the group of cyclic undecapeptides with anti-inflammatory, immunosuppressive, antifungal and antiparasitic properties (Sallam et al., 2005). It is used extensively in the prevention and treatment of graft-versus-host reactions in bone marrow transplantation and for the prevention of rejection of kidney, heart and liver transplants. Cyclosporin A is a major secondary metabolite usually produced by an aerobic filamentous fungus, *Tolypocladium niveum* (Park et al., 2006). Although cyclosporin A was initially developed as an antifungal antibiotic (Deo et al., 1984), it is currently prescribed as one of the most important immunosuppressive drugs for the treatment of organ transplants, as well as patients with autoimmune diseases, including AIDS, owing to its superior T-cell specificity and low levels of myelotoxicity (Borel, 1986). It consists of 11 amino acids with a molecular weight of 1202.6 and occurs as a white solid with a melting point of 148 to 151°C (natural) and 149 to 150°C (synthetic). It is stable in solution at temperatures below 30°C but is sensitive to light, cold, and oxidiza-

tion (IARC, 1990).

The drug was first isolated from *T. inflatum* (Gams, 1971) and after that very few studies were planned to explore other sources for its production. Until now most of the research work related to cyclosporin A deals with *T. inflatum* and *Aspergillus terreus*. The present study was designed to explore *Trichoderma* species for this drug production.

Materials and Methods

Acquisition and maintenance of *T. harzianum*

Pure cultures *T. harzianum* (FCBP 140), *T. aureoviride* (FCBP 234), *T. reesei* (FCBP 271), *T. koningii* (FCBP 585) and *T. hemantum* (FCBP 907) were obtained from the First Fungal Culture Bank of Pakistan (FCBP) and maintained on Malt extract agar (MEA) medium and preserved at 4°C.

Seed inoculum preparation

Malt Yeast extract (MY) medium was selected as seed



medium for inoculum preparation of selected fungi. MY medium is composed of malt extract 2%, yeast extract 0.4% and initial pH was adjusted to 5.7 using 1.0 M HCl. 50 mL of MY medium was prepared and poured in Erlenmeyer flasks (250 mL capacity) plugged with cotton plugs. These prepared medium flasks were sterilized by autoclaving at 121°C and 15 lb/inch² pressure for 15-20 min. By using cork borer, 0.8 cm disk of five days old cultures on MEA of *Trichoderma* species were inoculated in sterilized medium in flasks. The inoculated flasks were incubated on orbital shaker at 200 rpm for 72 hours at 30°C (Borel et al., 1977). The inoculum of all the strains was prepared by the same procedure.

Cultivation

For cultivation 50 mL production medium specially design for cyclosporin A production, was prepared with following composition: Glucose, 5%; peptone, 1%; KH₂PO₄, 0.5%; KCL, 0.25% (w/v), at pH 5.3. According to the methodology of Agathos et al. (1986) 5 mL of seed inoculum from each isolate was introduced into 250 mL Erlenmeyer flasks containing 50 mL of production medium. The fermentation was continued at 28 ± 1°C for 10 days, at 200 rpm.

Cyclosporin A extraction

Harvested fermentation medium was mixed with 30 mL of butyl acetate and stirred at 200 rpm for 24 hours at 30°C. Organic layer was formed after 24 hours, which was separated by separating funnel and evaporated under vacuum till dryness. Dried extract was weighed and dissolved in 30 mL methanol.

Measurement of fungal biomass

The aqueous layer of cultivation medium containing fungal pellets was filtered for harvest of biomass. The filtration was performed by using Whatman filter paper No. 1. The fresh biomass was collected on the filter paper by using a conical funnel. Filters were slightly rinsed with water 1-2 times for the washing of media and weighed on digital weighing machine. The filters were dried in oven at 40°C and re-weighed after cooling. Following formula was used to calculate the dry biomass:

Fungal dry biomass = weight of filter paper - weight of dry filtrate.

Cyclosporin A analysis

Level of cyclosporin A in the crude extract was analyzed by high performance liquid chromatography (HPLC). HPLC was carried out using Hitachi system consisting of L-2100/2130 pump, L-2420 UV-VIS detector with a detection span from 190 to 900 nm.

Analysis was done using a C₁₈ column with a 5 μm particle size, and acetonitrile:methanol:water

(42.5:20:37.5, v/v) as mobile phase at a flow rate of 0.8 mL/min with UV detection at 215 nm.

Cyclosporin A confirmation

For initial confirmation of cyclosporin A, Sandimmun Neoral® capsule (Novartis) containing 100 mg of cyclosporin was used. For this purpose 100 μL from capsule was taken with the help of micropipette and dissolved in 10 mL of methanol. Then, HPLC of this solution was carried out according to the conditions mentioned above.

The retention time and peak area of the samples were compared for final confirmation and quantification, with the external standard cyclosporin A 98.5% pure (Sigma-Aldrich, Fluka). The standard was analyzed under same conditions as described.

Estimation of cyclosporin A

The cyclosporin A level in crude extracts was determined by the following formula: % cyclosporin A by weight =

$$\frac{A_s W_r V_s}{A_r W_s V_r} \times \text{purity of reference}$$

Where, A_s is area of sample peak; A_r is area of reference peak; W_r is weight of reference material in grams; W_s is weight of sample in grams; V_s is volume of sample; V_r is volume of reference material.

The area of sample peaks and of reference peak was calculated from the chromatograms obtained by HPLC program LaChrom Elite.

Mass spectroscopy

The ESI/MS spectrum was obtained from single-quadrupole mass spectrometer. Sample was introduced using a silica capillary at a flow rate of 4.0 μL/min. The nebulizer gas was optimized and set at a rate of 1.6 L/min, and an electrospray potential of 4200 V was applied in the interface sprayer. A curtain gas of ultrapure nitrogen was pumped into the interface at the rate of 1.2 L/min.

Results and Discussion

Trichoderma as antagonists controlling wide range of pathogens are well documented (Ajith and Lakshmi-devi, 2010). As cyclosporin A is also known for its anti-fungal properties, five species of *Trichoderma* were checked for their potential to produce this drug. The drug extracted in butyl acetate was confirmed as cyclosporin A by comparing chromatograms with Sandimmun Neoral® capsules containing cyclosporin A as active ingredient and pure cyclosporin A. Sandimmun Neoral® capsules showed a clear peak at 2.77 whereas peak at 2.8 was recorded by authentic compound (Figure 1, Figure 2). In a previous study, Sallam et al.




Figure 1: HPLC chromatogram of Sandimmun Neoral^a capsule extracted at 215 nm showing cyclosporin A peak at 2.77




Figure 2: HPLC chromatogram of Sandimmun Neoral^a capsule extracted at 215 nm. Showing cyclosporin 'A' peak at 2.768




Figure 3: HPLC chromatograms of *T. harzianum* extracted at 215 nm. Peaks appeared at 2.779 in chromatogram identified as cyclosporin A

(2003) recorded cyclosporin A peak at 3.03 which was produced by *Aspergillus terreus* following almost similar experimental plan as is described in this study.

Among five tested *Trichoderma* species only *T. harzianum* (FCBP 140) isolated from air showed a comparable peak at 2.78 (Figure 3). The ESI-MS analysis showed the ion [CsA + H]⁺ at m/z 1203. The estimated yield of cyclosporine A was 44.1 µg/mL with a 0.26 g/50 mL of fungal biomass. No such peak was recorded in other tested species. However, this quantity was found lower to that recorded by earlier workers in the most studied fungal strain i.e., *T. inflatum*. As Dreyfuss et al. (1976) and Agathos et al. (1986) found a very good yield of 105.5 mg/L and Balakrishnan and Pandey (1996) showed that *T. inflatum* can produce higher levels of cyclosporin A up to 183 mg/L.

Beside *T. inflatum* so far this drug has been reported to be produced from *Aspergillus terreus*, *Fusarium solani* and *Fusarium oxysporum* (Sallam et al., 2003). This study confirms *T. harzianum* as a new fungal source of cyclosporine A production.

References

- Agathos SN, Marshall JW, Moraiti C, Parekh R, Modhosingh C. Physiological and genetic factors for process development of cyclosporin fermentation. *J Ind Microbiol.* 1986; 1: 39-48.
- Ajith PS, Lakshmidevi N. Effect of volatile and non-volatile compounds from *Trichoderma* spp. against *Colletotrichum capsici* incitant of Anthracnose on Bell peppers. *Nat Sci.* 2010; 8: 265-69.
- Borel JF, Feurer C, Magnee C, Stahelin H. Effects of the new anti-lymphocytic peptide cyclosporin A in animals. *Immunology* 1977; 32: 1017-25.
- Borel JF. Cyclosporin and its future In: Cyclosporin. Borel JF (ed). *Progress in Allergy.* Vol. 38. Switzerland, Karger, Basel, 1986, pp 9-18.
- Balakrishnan K, Pandey A. Growth and Cyclosporin A production by an indigenously isolated strain of *Tolypocladium inflatum*. *Folia Microbiol.* 1996; 41: 401-06.
- Deo YM, Gaucher GM. Semi-continuous and continuous production of penicillin G by *Penicillium chrysogenum* cells immobilized in κ-carrageenan beads. *Biotechnol Bioeng.* 1984; 26: 285-95.
- Dreyfuss M, Harri E, Hofmann H, Kobel H, Pache W, Tschertter H. Cyclosporin A and C: New metabolites from *Trichoderma polysporum*. *Eur J Appl Microbiol Biotechnol.* 1976; 3: 125-33.
- Gams W. *Tolypocladium*. Eine Hyphomycetengattung mit geschwollenen Phialiden. *Persoonia* 1971; 6: 185-91.
- IARC. *Pharmaceutical Drugs.* IARC Monographs on the evaluation of carcinogenic risk of chemicals to humans. International Agency for Research on Cancer, 1990, pp 415.
- Park NS, Park HJ, Han KB, Kim ES. Heterologous expression of novel cytochrome P450 hydroxylase genes from *Sebekia benihana*. *J Microbiol Biotechnol.* 2006; 16: 295-98.
- Sallam LA, El-Refai AM, Hamdi AH, El-Minofi HA, Abdel-Salam IS. Studies on the applications of immobilization technique for the production of cyclosporin A by a local strain of *Aspergillus terreus*. *J Gen Appl Microbiol.* 2005; 51: 143-49.
- Sallam LAR, El-Refai AMH, Hamdy AHA, El-Minofi HA, Abdel-Salam IS. Role of some fermentation parameters on cyclosporin A production by a new isolate of *A. terreus*. *J Gen Appl Microbiol.* 2003; 49: 321-28.

Author Info

Tehmina Anjum (Principal contact)
e-mail: tehminaanjum@yahoo.com