



BJP

Bangladesh Journal of Pharmacology

Research Article

**Evaluation of possible mechanisms
of three plants for blood glucose
control in diabetes**

Evaluation of possible mechanisms of three plants for blood glucose control in diabetes

Damayanthi Dalu¹ and Satyavati Dhulipala²

¹Department of Pharmacology, Netaji Institute of Pharmaceutical Science, Toopranpet, Choutuppal, Nalgonda 508 252, Telangana, India; ²Satyavati Dhulipala, Department of Pharmacology, Brilliant College of Pharmacy, Abdullapurmet, Hayatnagar, Ranga Reddy, Hyderabad 501 505, Telangana, India.

Article Info

Received: 8 September 2015
Accepted: 8 November 2015
Available Online: 24 January 2016
DOI: 10.3329/bjp.v11i1.24932

Cite this article:

Dalu D, Dhulipala S. Evaluation of possible mechanisms of three plants for blood glucose control in diabetes. Bangladesh J Pharmacol. 2016; 11: 224-30.

Abstract

This study was conducted to provide the evidence for the mechanism of anti-diabetic activity of *Cocculus orbiculatus*, *Leea indica* and *Ventilago maderaspatana*. This was accomplished by employing methods like uptake of glucose, glycogen synthesis and inhibition of α -glucosidase. For uptake of glucose, diaphragms were dissected out in Tyrode solution with 2% glucose and assayed for glucose content. In glycogen synthesis methodology liver, skeletal muscle and cardiac muscles were isolated, homogenized and glycogen content was analyzed. In α -glucosidase enzyme inhibition procedure involved estimation of α -glucosidase enzyme inhibition. All the three plant extracts exhibited significant ($p < 0.05$ - $p < 0.01$) anti-diabetic activity by increasing glucose uptake, glycogen synthesis and inhibiting α -glucosidase enzyme. Among the three plants, *V. maderaspatana* (500 mg/kg) exhibited higher glucose uptake, glycogen content and α -glucosidase inhibition activity (IC_{50} 145 μ g/mL). The present experimental results evidenced the anti-diabetic activity of three plants by all the three mechanisms.

Introduction

Diabetes mellitus is characterized by impaired production of insulin and/or diminished stimulation of insulin sensitive peripheral tissues associated with a marked decrease in glucose uptake and metabolism in response to insulin. The defective glucose transport system plays a significant role in the pathogenesis of peripheral insulin resistance. Glucose uptake in target tissues is a crucial step in maintaining glucose homeostasis and in lowering the postprandial glucose load (Shulman, 2000). Direct stimulation of glucose transport and metabolism in muscle and fat cells lead to enhanced glucose utilization. To attenuate the glucose uptake by peripheral cells, biguanides are employed. Cellular assays are utilized to investigate the mechanism of action of natural compounds using isolated rat diaphragms. It is highly preferred to explore modern anti-diabetic agents from natural sources that stimulate glucose uptake/

disposal by peripheral tissues such as adipose tissue or muscle cells.

The pivotal enzyme for carbohydrate digestion is α -glucosidase. This is a therapeutic target for the modulation of postprandial hyperglycemia, the earliest abnormality that occurs in NIDDM (Kim et al., 2005). Dietary carbohydrates are the major source for blood glucose. These carbohydrates are hydrolyzed by α -glucosidase, so as to be absorbed by small intestine. Therefore, the most effective treatment is to inhibit the activity of α -glucosidase (Krentz and Bailey, 2005). α -Glucosidase inhibitors such as acarbose, miglitol and voglibose reduce postprandial hyperglycemia by inhibiting the activity of carbohydrate digesting enzymes and delaying glucose absorption.

Previously, we have reported anti-diabetic, anti-hyperlipidemic and anti-oxidant activity of three medicinal



plants *Cocculus orbiculatus*, *Leea indica* and *Ventilago maderaspatana* in the treatment of diabetes (Damayanthi and Satyavati, 2015); Damayanthi et al., 2014; Damayanthi and Satyavati, 2015). But, till date there are no scientific evaluation reports available to support the mechanisms responsible for anti-diabetic activity. Therefore, the present investigation was aimed to ascertain *in vivo* anti-diabetic activity by methods such as glucose uptake activity using isolated rat diaphragm and glycogen synthesis in liver, skeletal muscle and cardiac muscle and *in vitro* anti-diabetic activity by inhibiting α -glucosidase enzyme.

Materials and Methods

Plant materials

Aerial parts of *C. orbiculatus* were collected from Tirumala forest area, Tirupathi. *L. indica* leaves were procured from Karthikavanam forest area, Dhulapally, Hyderabad. *V. maderaspatana* roots were obtained from Tirumala forest area, Tirupathi. The plants were authenticated by Prof. Madhava Chetty, Department of Botany, Sri Venkateshwara University, Tirupathi, India.

Chemicals

Streptozotocin was procured from Sigma-Aldrich. Glucose estimation kit was obtained from Erba Diagnostics, Mannheim. Glibenclamide (Oglucon) was purchased from Alpha pharmaceuticals, Apollo pharmacy, Bathala-palli, Ananthapur. Insulin (Novo Nordisk) was obtained from Alpha pharmaceuticals, Bathalapalli, Ananthapur. Glucose was purchased from Sd Fine Chemicals, India.

Preparation of plant extract

Aerial parts of *C. orbiculatus* were finely powdered, packed in soxhlet apparatus and then extracted with hydroalcohol (60:40). Leaves of *L. indica* were air dried at room temperature, coarsely powdered extracted by maceration with hydroalcohol (3:1). *V. maderaspatana* roots were finely powered and extracted by using soxhlet apparatus with hydroalcohol solvent (60:40). Percentage yield of the plants *C. orbiculatus*, *L. indica*, *V. maderaspatana* was found to be 16.7, 25.6 and 15.8% respectively.

Preliminary phytochemical analysis

All the three plant extracts were subjected to preliminary phytochemical analysis to determine the phyto-constituents employing standard tests (Harbone, 1998).

Animals

Wistar albino rats weighing about 200-250 g were procured from Raghavendra enterprises, Bangalore. The animals were acclimatized (2 weeks); housed under standard laboratory conditions (temperature $23 \pm 2^\circ\text{C}$),

humidity 55-70% and fed with commercial diet Durga feeds, Bangalore.

α -glucosidase inhibitory assay

This assay was assigned to investigate the *in vitro* inhibitory activity of three plant extracts on α -glucosidase enzyme. α -Glucosidase (100 μL of 1 U/mL) was mixed with phosphate buffer (100 μL , pH 7.0) containing 100 μL of three plant extracts (25-1600 μL) or standard drug acarbose (0.1-3.2 $\mu\text{g}/\text{mL}$). This mixture is incubated at 37°C for 60 min in maltose solution. Later the mixture is kept in boiling water for 2 min and cooled. The boiling stops the α -glucosidase action on maltose. Glucose reagent (2 mL) was added and absorbance is measured at 540 nm to estimate the amount of liberated glucose by the action of α -glucosidase (Kuppusamy et al., 2011)

Glucose uptake in normal and streptozotocin-induced diabetic rats

The glucose uptake using rat hemidiaphragm was estimated according to the method reported elsewhere (Walass and Walass, 1952; Chattopadhyay et al., 1992), but with some modifications. After 18 hours fasting; rats were killed by decapitation. Diaphragms were dissected out quickly with minimal trauma and divided into two halves. Hemidiaphragms were then rinsed in cold Tyrode solution (without glucose) to remove blood clots. Then these were weighed and placed in test tubes. The volumes in all test tubes were made equal by adding distilled water. The test tubes were incubated for 30 min at 37°C in an atmosphere of 100% oxygen and were shaken at 140 cycles/min. Hemidiaphragms were taken out. Glucose content of the incubated medium before and after incubation was measured. This was carried out by employing Erba Diagnostic Mannheim kit using GOD-POD method (Barham and Trinder, 1972) and Erba Mannheim Chem-7 semiautoanalyzer. Glucose uptake was calculated as the difference between initial and final glucose content. Glucose uptake was expressed as mg/g of tissue per 30 min of incubation. Rats were divided into 12 groups; six in normal group and six in diabetic control group of five rats each. In normal group: Group I (normal rats; received 2 mL Tyrode solution with 2% glucose); Group II (received Tyrode solution and 0.6 mL of 0.4 IU/mL insulin); Group III (administered Tyrode solution and standard drug, metformin- 2 mL of 0.1%); Group IV (received Tyrode solution and 300 mg/kg of *C. orbiculatus*); Group V (received Tyrode solution- 400 mg/kg of *L. indica*); Group VI (administered Tyrode solution and 500 mg/kg *V. maderaspatana*).

In diabetic control group: Group VII (diabetic control group, untreated); Group VIII (received Tyrode solution and 0.6 mL of 0.4 IU/mL insulin); Group IX (administered Tyrode solution and standard drug, metformin- 2 mL of 0.1%); Group X (received Tyrode solution and 300 mg/kg *C. orbiculatus*);

Group XI (received Tyrode solution and 400 mg/kg *L. indica*); Group XII (administered Tyrode solution and 500 mg/kg *V. maderaspatana*)

Glycogen estimation in normal and streptozotocin-induced diabetic rats

The glycogen content in liver, skeletal muscle and cardiac muscle was estimated by Carroll et al., (1956). Rats were divided into ten groups: Five in normal group and five in diabetic control group of five rats each.

Normal group: Group I (normal rats; received 1% sodium carboxymethyl cellulose); Group II (received standard drug, glibenclamide 10 mg/kg); Group III (administered 300 mg/kg of *C. orbiculatus*); Group IV (administered 400 mg/kg of *L. indica*) Group V (administered 500 mg/kg *V. maderaspatana*)

Diabetic control group: Group VI (diabetic control group, untreated); Group VII (diabetic control received standard drug, glibenclamide 10 mg/kg); Group III (diabetic control administered 300 mg/kg of *C. orbiculatus*); Group IV (diabetic control administered 400 mg/kg of *L. indica*); Group V (diabetic control administered 500 mg/kg of *V. maderaspatana*)

After 18 hours of fasting, plant extracts were administered to different groups. Two hours later they were sacrificed by decapitation. The liver, skeletal muscle and cardiac muscle were isolated, weighed and homogenized using 10 mL of 4% trichloroacetic acid and centrifuged for 10 min. Supernatant was decanted and precipitate is discarded. To 2 mL of supernatant 4 mL of

anthrone reagent was added. Later test tubes were allowed to cool for 30 min. Absorbance was measured at 620 nm using spectrophotometer. Glycogen content was expressed as milligram for 100 g of tissue.

Glycogen content = $DU \times 0.2 \times \text{volume of the extract} \times 1000 \text{ DS} \times \text{weight of the tissue}$

DU= Absorbance of the sample; DS= Absorbance of the standard

Statistical analysis

The experimental results were presented as mean \pm standard error mean (SEM). Statistical analysis was performed by graphpad instat version 3.2. Probability value of analysis $p < 0.01$ and $p < 0.05$ was considered to be statistically significant.

Results

Preliminary phytochemical analysis

Phytochemical analysis of *C. orbiculatus* exhibited positive results for alkaloids, glycosides, carbohydrates, flavonoids, saponins, tannins, terpenoids, polyphenols and starches (Table I). Phytochemical analysis of *L. indica* exhibited the presence of alkaloids, terpenoids, carbohydrates, flavonoids, tannins and saponins. Preliminary phytochemical analysis of *V. maderaspatana* revealed the presence of alkaloids, glycosides, emodin, cardiac glycosides, carbohydrates, flavonoids, tannins and saponins. The constituents like aporphine

Table I

Phytochemical screening of *C. orbiculatus*, *L. indica* and *V. maderaspatana*

Phytoconstituents	<i>C. orbiculatus</i>	<i>L. indica</i>	<i>V. maderaspatana</i>
Alkaloids	++	++	++
Glycosides	++	--	++
Carbohydrates	++	++	++
Terpenoids	++	++	++
Flavonoids	++	++	++
Saponins	++	++	++
Tannins	++	++	++
Starches	++	--	--
Polyphenols	++	--	--
Steroids	--	++	--
Gallic acid	--	++	--
β -Sitosterol	--	++	--
Cardiac glycosides	--	++	++
Emodin	--	--	++
Dam-karrer test	--	--	++
Juglone test	--	--	++

Note: Present (++); Absent (--)

Table II			
Effect of <i>C. orbiculatus</i> , <i>L. indica</i> and <i>V. maderaspatana</i> on inhibition of α -glucosidase enzyme			
Treatment	Concentration ($\mu\text{g/mL}$)	% Inhibition	IC ₅₀ ($\mu\text{g/mL}$)
<i>C. orbiculatus</i>	25	8.3 \pm 0.5	325.1
	50	17.1 \pm 0.5	
	100	23.2 \pm 0.4	
	200	37.1 \pm 0.5	
	400	55.3 \pm 1.7	
	800	67.3 \pm 1.2	
	1000	85.7 \pm 1.4	
<i>L. indica</i>	25	10.3 \pm 0.4	265.3
	50	20.5 \pm 0.7	
	100	33.7 \pm 0.5	
	200	45.3 \pm 0.8	
	400	62.2 \pm 0.9	
	800	79.5 \pm 1.3	
	1000	90.3 \pm 1.8	
<i>V. maderaspatana</i>	25	15.1 \pm 0.3	145.0
	50	29.3 \pm 0.4	
	100	43.7 \pm 0.7	
	200	56.3 \pm 0.5	
	400	75.3 \pm 1.0	
	800	88.9 \pm 1.1	
	1000	95.8 \pm 1.3	
Acarbose	0.1	30.7 \pm 0.4	0.2
	0.2	47.3 \pm 0.7	
	0.4	59.2 \pm 1.1	
	0.8	73.1 \pm 2.1	
	1.6	81.2 \pm 1.9	
	3.2	96.2 \pm 1.7	

and berberine, ursolic acid, gallic acid, β -sitosterol, emodin and physcion were found by the analysis of *C. orbiculatus*, *L. indica* and *V. maderaspatana*.

α -Glucosidase inhibitory activity

C. orbiculatus, *L. indica* and *V. maderaspatana* hydroalcoholic extracts exhibited 8.3, 10.3 and 15.1% inhibition of α -glucosidase activity at 25 $\mu\text{g/mL}$ and 85.7, 90.3 and 95.8% inhibition at 1600 $\mu\text{g/mL}$ respectively (Table II). The IC₅₀ values were 325, 265 and 145 $\mu\text{g/mL}$. IC₅₀ value of acarbose was found to be 0.2 $\mu\text{g/mL}$.

Effect on peripheral glucose uptake

Table III show glucose uptake in an isolated rat hemidiaphragm muscle of normal and diabetic animals. Addition of *C. orbiculatus*, *L. indica* and *V. maderaspatana* hydroalcoholic extracts elicited significant increase in glucose uptake by the rat hemidiaphragm in normal animals. *V. maderaspatana* seemed to be more effective

in enhancing peripheral glucose uptake than *L. indica*, *C. orbiculatus* and metformin. The glucose uptake by rat hemidiaphragm was significantly higher in all the groups when compared to control group. Diabetic control animals exhibited significant increase in glucose uptake of *C. orbiculatus*, *L. indica* and *V. maderaspatana* (37.7, 40.8 and 48.0% respectively). Glucose uptake of *V. maderaspatana* was significantly higher when compared to *L. indica* and *C. orbiculatus*.

Effect on glycogen content in liver, skeletal muscle and cardiac muscle in normal animals

Hydroalcoholic extracts of *C. orbiculatus*, *L. indica* and *V. maderaspatana* showed significantly increased glycogen content in liver (Table IV). Increase in glycogen content was more for *V. maderaspatana* than *L. indica*, *C. orbiculatus* and glibenclamide. Skeletal muscle glycogen content was increased more for all the three plant extracts. Increase was greater for *V. maderaspatana* than *L. indica*, *C. orbiculatus* and glibenclamide. Significant

Table III

Effect of *C. orbiculatus*, *L. indica* and *V. maderaspatana* on glucose uptake in isolated rat hemidiaphragm (mg/g/30 min)

Group	Treatment (mg/kg)	Glucose uptake in normal rats	Glucose uptake in diabetic rats
I	Normal	5.4 ± 0.15	4.7 ± 0.2
II	Insulin (0.4 IU)	12.7 ± 0.6 ^b (57.5%)	11.5 ± 0.8 ^b (59.2%)
III	Metformin (0.1%)	9.6 ± 0.2 ^b (43.5%)	8.3 ± 0.9 ^b (43.6%)
IV	<i>C. orbiculatus</i> (300)	8.4 ± 0.3 ^b (35.7%)	7.5 ± 0.4 ^b (37.7%)
V	<i>L. indica</i> (400)	9.1 ± 0.5 ^b (40.7%)	7.9 ± 0.2 ^b (40.8%)
VI	<i>V. maderaspatana</i> (500)	11.3 ± 0.7 ^b (52.4%)	9.0 ± 0.1 ^b (48.0%)

IU: International units; Mean ± SEM; n = 5; Groups II - VI were compared with Group I. ^b(p<0.01)

Table IV

Effect of *C. orbiculatus*, *L. indica* and *V. maderaspatana* on glycogen content in liver, skeletal muscle and cardiac muscle (mg/100 g)

Group	Treatment	Glycogen concentration in normal rats			Glycogen concentration in diabetic rats		
		Liver	Skeletal muscle	Cardiac muscle	Liver	Skeletal muscle	Cardiac muscle
I	Normal	136.5 ± 2.3	30 ± 1.2	29 ± 1.4	116 ± 1.2	22 ± 1.4	21 ± 1.0
II	Glibenclamide (10)	153.0 ± 2.0 ^b	39.5 ± 0.8 ^b	40.5 ± 1.3 ^b	143 ± 1.0 ^b	33.7 ± 0.6 ^b	37 ± 0.7 ^b
III	<i>C. orbiculatus</i> (300)	142.2 ± 2.0 ^a	35 ± 1.1 ^a	32.7 ± 0.4 ^a	126 ± 0.8 ^b	26 ± 0.8 ^a	26 ± 0.7 ^b
IV	<i>L. indica</i> (400)	150.7 ± 2.0 ^b	38 ± 1.9 ^b	33.7 ± 0.6 ^b	131.3 ± 0.9 ^b	30 ± 0.8 ^b	33 ± 1.1 ^b
V	<i>V. maderaspatana</i> (500)	162.5 ± 2.9 ^b	40 ± 0.9 ^b	41.0 ± 0.7 ^b	139 ± 1.0 ^b	32.7 ± 1.0 ^b	36.7 ± 1.3 ^b

Mean ± SEM; n = 5; Groups II - VI were compared with Group I. ^a(p<0.05); ^b(p<0.01)

increase in cardiac muscle glycogen content was observed for *V. maderaspatana* than *L. india* and *C. orbiculatus*. *V. maderaspatana* exhibited increased cardiac muscle glycogen content than *L. india*, *C. orbiculatus* and glibenclamide.

Effect on glycogen content in liver, skeletal muscle and cardiac muscle in diabetic animals

Three plants have elicited significantly increased liver glycogen content. *V. maderaspatana* manifested higher liver glycogen content than *L. indica* and *C. orbiculatus*. Significantly increased skeletal muscle glycogen content was observed for *V. maderaspatana*, *L. indica* and *C. orbiculatus*. Glycogen content was more in *V. maderaspatana* than *L. indica* and *C. orbiculatus*. Cardiac muscle glycogen content increase was observed for all the three plants. Among these *V. maderaspatana* exhibited increased glycogen content than *L. indica* and *C. orbiculatus*.

Discussion

In current study, we investigated the mechanism for

anti-diabetic activity of three plants (*C. orbiculatus*, *L. indica* and *V. maderaspatana*) by employing glucose uptake, glycogen synthesis and α -glucosidase enzyme inhibition methods. Three plants showed significant uptake of glucose and are more or less effective than insulin. *V. maderaspatana* has evidenced greater uptake of glucose than *L. india*, *C. orbiculatus*, metformin and comparatively equal to insulin. Glibenclamide showed increased glycogen content in muscle cells. Therapy with the three plants evidenced increased glucose uptake and thereby glycogen storage in liver, skeletal muscle and cardiac muscle. *V. maderaspatana* increased glycogen content greater than *L. india* and *C. orbiculatus*. The three plants also evidenced α -glucosidase inhibitory activity among which *V. maderaspatana* exhibited greater activity.

The impaired glucose uptake is linked with decrease in the translocation of GLUT 4 and is the major cause of insulin resistance. Metformin and insulin stimulate glucose uptake in muscle cells by increasing GLUT 4 (Klip and Leiter, 1990). Berberine the reported active constituent of *C. orbiculatus* has been investigated to activate AMPK in skeletal muscle and adipose tissue that lead to

increased glucose uptake (Yin et al., 2008; Cheng et al., 2006; Kim et al., 2007; Lee et al., 2006; Zhou et al., 2007; Wang et al., 2004; Ko et al., 2005). Similarly gallic acid the constituent of *L. india* has been implicated to mediate insulin stimulated glucose transport in muscle and adipocytes (Vishnu Prasad et al., 2010). Finally VMHAE possess emodin that mediated Glut 1 and Glut 4 expression to enhance glucose uptake in skeletal muscles and adipocytes (Ying Yang et al., 2007). Glibenclamide increased glycogen content in muscle but adverse effects like hypoglycemia and weight gain limits the use. Hence drugs that attenuate glycogen content without adverse effects were desirable for long term anti-diabetic therapy. The three plants evidenced enhanced glucose uptake and glycogen storage in liver, skeletal muscle and cardiac muscle. The increased glycogen levels by the three plants might be due to the berberine of *C. orbiculatus* that evidenced glucose utilization in Hep G₂ cells and 3T3-L1 adipocytes increasing glycogen content (Yin J et al., 2002; Zhou et al., 2003; Zhou et al., 2003). Gallic acid, the constituent of *L. india* by insulin secretagogue action produced increased glycogen levels (Punithavathi et al., 2011). Emodin of *V. maderaspatana*, the natural PPAR γ activator, up-regulated PPAR γ and increased glycogen content (Yonemitsu et al., 2001). Acarbose, the α -glucosidase inhibitor has common side effects (flatulence and abdominal bloating). Herbal drugs devoid of side effects are desired to improve compliance for diabetic patients. The constituents like berberine, β -sitosterol and emodin of the plants have reported α -glucosidase inhibitory activity (Pan et al., 2003; Sunil Kumar et al., 2013; Yang et al., 2014). Thus active constituents of the experimental study might be responsible for the elicited α -glucosidase inhibitory activity of *C. orbiculatus*, *L. indica* and *V. maderaspatana*.

Conclusion

The experimental reports evidenced *C. orbiculatus*, *L. indica* and *V. maderaspatana* plants act through multiple mechanisms like increased glucose uptake, glycogen synthesis in muscle cells and α -glucosidase inhibition to control blood glucose in diabetes. *V. maderaspatana* elicited higher antidiabetic activity compared to *L. indica* and *C. orbiculatus*.

Financial Support

Self-funded

Ethical Issue

Experiments were conducted in accordance with the guidelines of CPCSEA REGD No. 878/ac/05/CPCSEA/21/2015. The study protocol was approved by Institutional Animal ethics committee.

Conflict of Interest

Authors declare no competing interest

Acknowledgements

The authors express deep gratitude to N. Raja Kumar, Y. Ganesh Kumar and S. Nagarjuna whose valuable contributions lead to successful completion of the work. The authors also thank the College Raghavendra Institute of Pharmaceutical Education and Research for providing necessary facilities.

References

- Barham D, Trinder P. An improved colour reagent for the determination of blood glucose by the oxidase system. *Analyst* 1972; 97: 142-45.
- Carroll NV, Longely RW, Rose JH. The determination of glycogen in liver and muscle by use of anthrone reagent. *J Biol Chem.* 1956; 220: 583-93.
- Chattopadhyay RR, Sarkar SK, Ganguly S, Banerjee RN, Basu TK. Effect of leaves of *Vinca rosea* Linn. on glucose utilization and glycogen deposition by isolated rat hemidiaphragm. *Indian J Physiol Pharmacol.* 1992; 36: 137-38.
- Cheng Z, Pang T, Gu M, Gao AH, Xie CM, Li JY, Nan FJ, Li J. Berberine-stimulated glucose uptake in L6 myotubes involves both AMPK and p38 MAPK. *Biochim Biophys Acta.* 2006; 1760: 1682-89.
- Damayanthi D, Satyavati D, Suresh A. Antihyperglycemic and hypolipidemic activity of *Lea indica*. *Int J Bioassays.* 2014; 3: 3155-59.
- Damayanthi D, Satyavati D. Anti-diabetic and anti-hyperlipidemic activity of alcoholic and hydroalcoholic extracts of *Cocculus orbiculatus* in streptozotocin induced diabetic rats. *Am J Phytomed Clin Ther.* 2015; 3: 276-86.
- Damayanthi D, Satyavati D. Anti-diabetic, anti-hyperlipidemic and anti-oxidant properties of roots of *Ventilago maderaspatana* Gaertn. on streptozotocin-induced diabetic rats. *IOSR: J Pharm Biol Sci.* 2015; 10: 50-59.
- Harbone JB. *Phytochemical methods: A guide to modern techniques of plant analysis.* 3rd ed. New York, Chapman and Hall Publisher, Springer, 1998, p 302.
- Kim YM, Jeong YK, Wang MH, Lee WY, Rhee HI. Inhibitory effect of pine extract on α -glucosidase activity and postprandial hyperglycemia. *Nutrition* 2005; 21: 756-61.
- Kim SH, Shin EJ, Kim ED, Bayaraa T, Frost SC, Hyun CK. Berberine activates GLUT1-mediated glucose uptake in 3T3-L1 adipocyte. *Biol Pharm Bull.* 2007; 30: 2120-25.
- Klip A, Leiter L.A. Cellular mechanism of action of metformin. *Diabetes Care.* 1990; 13: 696-704.
- Ko BS, Choi SB, Park SK, Jang JS, Kim YE, Park S. Insulin sensitizing and insulinotropic action of berberine from *Cortidis rhizome*. *Biol Pharm Bull.* 2005; 28: 1431-37.
- Krentz AJ, Bailey CJ. Oral antidiabetic agents: Current role in type 2 diabetes mellitus. *Drugs* 2005; 65: 385-411.

- Kuppusamy A, Muthusamy U, Andichetiar Thirumalaisamy S, Varadharajan S, Ramasamy K, Ramanathan S. *In vitro* (α -glucosidase and α -amylase inhibition) and *in vivo* antidiabetic property of phytic acid (IP 6) in streptozotocin-nicotinamide induced type 2 diabetes mellitus (NIDDM) in rats. *J Complement Integrat Med*. 2011; 8.
- Lee YS, Kim WS, Kim KH, Yoon MJ, Cho HJ, Shen Y, Ye JM, Lee CH, Oh WK, Kim CT. Berberine a natural plant product, activates AMP-activated protein kinase with beneficial metabolic effects in diabetic and insulin-resistant states. *Diabetes* 2006; 55: 2256-64.
- Pan GY, Huang ZJ, Wang GJ, Fawcett JP, Liu XD, Zhao XC, Sun JG, Xie YY. The anti-hyperglycaemic activity of berberine arises from a decrease of glucose absorption. *Planta Med*. 2003; 69: 632-36.
- Punithavathi VR, Stanely Mainzen Prince P, Kumar MR, Selva Kumara CJ. Protective effects of gallic acid on hepatic lipid peroxide metabolism, glycoprotein components and lipids in streptozotocin-induced type II diabetic Wistar rats. *J Biochem Molecular Toxic*. 2011; 25: 68-76.
- Shulman GI. Cellular mechanisms of insulin resistance. *J Clin Invest*. 2000; 106: 171-76.
- Sunil Kumar, Vipin Kumar, Om Prakash. Enzymes inhibition and Anti-diabetic effect of isolated constituents from *Dillenia indica*. *Biomed Res Int*. 2013; 2013: 1155-62.
- Vishnu Prasad CN, Anjana T, Banerji A, Gopalakrishnapillai A. Gallic acid induces GLUT 4 translocation and glucose uptake activity in 3T3 cells. *FEBS Lett*. 2010; 584: 531-36.
- Walass E, Walass O. Effect of insulin on rat diaphragm under anaerobic conditions. *J Biol Chem*. 1952; 195: 367-73.
- Wang SH, Wang WJ, Wang XF, Chen W. Effect of *Astragalus polysaccharides* and berberine on carbohydrate metabolism and cell differentiation in 3T3-L1 adipocytes. *Chin J Integr Tradit Chin West Med*. 2004; 24: 926-28.
- Yang D, Zhao J, Liu S, Song F, Liu ZQ. The screening of potential α -glucosidase inhibitors from the *Polygonum multiflorum* extract using ultra-filtration combined with liquid chromatography-tandem mass spectrometry. *Analyt Methods*. 2014; 6: 3353-59.
- Yin J, Hu R, Chen M, Tang J, Li F, Yang Y, Chen J. Effects of berberine on glucose metabolism *in vitro*. *Metabolism* 2002; 51: 1439-43.
- Yang Y, Shang W, Zhou L, Jiang B, Jin H, Chen M. Emodin with PPAR γ ligand-binding activity promotes adipocyte differentiation and increases glucose uptake in 3T3-L1 cells. *Biochem Biophys Res Comm*. 2007; 353: 225-30.
- Yonemitsu S, Nishimura H, Shintani M, Yoshihiro O, Tatsuya H, Kiminori H, Inoue G, Kazuwa N. Troglitazone induces GLUT 4 translocation in L6 myotubes. *Diabetes* 2001; 50: 1093-101.
- Yin J, Gao Z, Liu D, Liu Z, Ye J. Berberine improves glucose metabolism through induction of glycolysis. *Am J Physiol Endocrinol Metab*. 2008; 294: E148-56.
- Zhou LB, Chen MD, Song HD, Yang Y, Wang X, Tang JF, Li FY, Xu MY, Chen JL. Effect of berberine on glucose transport in adipocytes and its mechanism. *Chin J Endocrinol Metab*. 2003; 19: 479-82.
- Zhou LB, Chen MD, Wang X, Song HD, Yang Y, Tang JF, Li FY, Xu MY, Chen JL. Effect of berberine on the differentiation of adipocytes. *Zhonghua Yi Xue Za Zhi*. 2003; 83: 338-40.
- Zhou L, Yang Y, Wang X, Liu S, Shang W, Yuan G, Li F, Tang J, Chen M, Chen J. Berberine stimulates glucose transport through a mechanism distinct from insulin. *Metabolism* 2007; 56: 405-12.

Author Info

Damayanthi Dalu (Principal contact)

e-mail: damayanthidalu3@gmail.com; Cell: +91 9700550851