


# Bangladesh Journal of Pharmacology

Volume: 12; Number 4; Year 2017



Cite this article as: Sezen Karaoglan E, Özgen U, Kazaz C. Phytochemical studies on *Origanum rotundifolium*. Bangladesh J Pharmacol. 2017; 12: 470-471.



## Letter to the Editor

### Phytochemical studies on *Origanum rotundifolium*

Sir,

The genus *Origanum* is one of the important genera of the Lamiaceae family. *Origanum* species have rich essential oil constituents and these plants are used as spices. They are also extensively used in the flavouring of food products and alcoholic drinks and perfumery for their spicy smell. *Origanum* species are used as antiseptic, stimulant, stomachic, expectorant, sudofiric and emmenagogic in the folk medicine (Aligiannis et al., 2001; Baser, 1978; Novak et al., 2000).

*Origanum rotundifolium* is a member of *Origanum* genus. This medicinal plant known in Anatolia as "Mercan kösk" is used as spices. Thymol, carvacrol, *p*-cymene and borneol were found to be as main constituents of the essential oil of *O. rotundifolium*. The essential oil of this plant shows various biological activities such as anti-oxidant, antibacterial properties. These activities have been mainly attributed to constituents such as thymol, carvacrol, borneol, terpinenchemo type and spathulenol (Baser et al., 1993; Gormez et al., 2016; Goze et al., 2009; Kotan et al., 2010).

In the present study, we report on the isolation and structure elucidation of 3 metabolites of *O. rotundifolium*.

The aerial parts of *O. rotundifolium* were collected from Barhal (Yusufeli, Artvin Province, 560-620 m, Turkey). A voucher specimen was deposited at the Herbarium of Ankara University, Faculty of Pharmacy (AEF25873).

The air-dried and powdered aerial parts (400 g) of *O. rotundifolium* were extracted three times with MeOH at 40°C (3 × 2L). After filtration, the methanol extracts were evaporated under vacuum to dryness. Methanol extract (71.6 g) was dissolved in water-methanol (9:1) and partitioned with chloroform and then ethyl acetate, which were separately concentrated and dried under reduced pressure to give 18.3 g and 4.0 g residues, respectively. The remaining aqueous phase was 31.0 g.

Chloroform extract (2.7 g) was separated via silica gel column chromatography eluting with *n*-hexane-ethyl acetate (100:0, 90:10 ... 50:50). Fr. 6-7 gave the compound **1** (45 mg).

Ethyl acetate extract was subjected to reversed phase silica gel column chromatography using water: methanol (90:10, 80:20.....,100) solvent systems. The fractions 4-10 (1.79 g) were further purified by consecutive column chromatography on silica gel (chloroform-methanol-water, 80:20:2, 70:30:3,...., 50:50:5) and sephadex LH-20 (MeOH), respectively. Fr. 8 gave compound **2** (44 mg).

The remaining aqueous phase was subjected to reversed phase silica gel column chromatography using water: methanol (90:10, 80:20.....,100) solvent systems. Fr. 19-29 were separated via silica gel column chromatography eluting with chloroform-methanol-water (80:20:2, 70:30:3,...., 50:50:5). Fr. 84-121 gave compound **3** (217 mg).

Their structures were identified by means of spectroscopic methods (1D- and 2D-NMR, EIMS).

#### Compound 1

EIMS  $m/z$  236 [M+Na-H]<sup>+</sup>, C<sub>14</sub>H<sub>30</sub>O<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz)δ: 3.66 (2H, t, J= 6.6 Hz, H-1), 1.28-1.60 (24H, H-2-13), 0.90 (3H, t, J= 6.8 Hz, H-14). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100 MHz) δ:22.7, 25.7, 29.4, 29.4, 29.6, 29.7, 29.7, 31.9, 32.82 (C-2-13), 14.1 (C-14).

#### Compound 2

C<sub>18</sub>H<sub>16</sub>O<sub>8</sub>,<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz)δ: 7.02 (1H, d, J= 1.8 Hz, H-2), 6.76 (1H, d, J= 8.4 Hz, H-5), 6.92 (1H, dd, J= 8.1, J= 2.2 Hz, H-6), 7.50 (1H, d, J= 16.0 Hz, H-7), 6.27 (1H, d, J= 16.0 Hz, H-8), 6.75 (1H, d, J= 1.8 Hz, H-2'), 6.67 (1H, d, J= 8.1 Hz, H-5'), 6.62 (1H, dd, J= 8.1 Hz, J= 1.8 Hz, H-6'), 3.09 (1H, dd, J= 14.1 Hz, J= 3.1 Hz, Ha-7'), 2.92 (1H, dd, J= 14.1 Hz, J= 9.7 Hz, Hb-7'), 5.08 (1H, dd, J= 9.7, J= 3.5 Hz, H-8'). <sup>13</sup>C-NMR (CDCl<sub>3</sub>, 100 MHz) δ: 126.8 (C-1), 114.4 (C-2), 145.5 (C-3), 148.2 (C-4), 115.3 (C-5), 121.7 (C-6), 145.5 (C-7), 113.9 (C-8), 167.9 (C-9), 129.9 (C-1'), 116.3 (C-2'), 144.8 (C-3'), 143.6 (C-4'), 115.0 (C-5'), 120.6 (C-6'), 37.6 (C-7'), 76.3 (C-8'), 176.1 (C-9').

#### Compound 3

EIMS  $m/z$  561 [M+Na]<sup>+</sup>, C<sub>27</sub>H<sub>22</sub>O<sub>12</sub>, <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz)δ: 7.10 (1H, d, J= 8.4 Hz, H-6), 7.95 (1H, d, J= 16.0 Hz, H-7), 6.24 (1H, d, J= 16.0 Hz, H-8), 5.11 (1H, dd, J= 6.2 Hz, H-10), 3.07 (bd, J= 13.0 Hz, Ha-11), 2.94 (dd, J= 13.0 Hz, J= 9.0 Hz, Hb-11), 6.87 (1H, s, H-13), 6.63 (d, J= 8.0 Hz, H-16), 6.57 (bd, J= 8.0 Hz, H-17), 4.26 (1H, d, J= 6.2 Hz, H-20), 5.87 (1H, d, J= 6.2 Hz, H-21), 6.85 (1H, s, H-23), 6.71-6.75 (3H, m, signaloverlap). <sup>13</sup>C-NMR






Figure 1: Compounds isolated from *O. rotundifolium*

(CDCl<sub>3</sub>, 100 MHz)  $\delta$ :123.6 (C-1), 129.0 (C-2), 147.5 (C-3), 145.3 (C-4), 116.6 (C-5), 119.9 (C-6), 142.7 (C-7), 116.0 (C-8), 167.8 (C-9), 75.9 (C-10).

At the end of the extraction and isolation processes of aerial parts of *O. rotundifolium* 71.6 g methanol extract was obtained. Chloroform, ethyl acetate and remaining aqueous phase were 18.3 g, 4 g and 31 g, respectively after fractionation. 1-Tetradecanol (Compound **1**, 45 mg) was isolated from the chloroform phase; rosmarinic acid (Compound **2**, 44 mg) from ethyl acetate phase; lithospermic acid (Compound **3**, 217 mg) from the aqueous phase (Figure 1). <sup>1</sup>H-NMR and <sup>13</sup>C-NMR data comply with the data given in the literature (Junges et al., 2000; Kelley et al., 1975; Kelley et al., 1976).

Earlier studies have shown that *O. rotundifolium* has anti-oxidant, antibacterial effects, including essential oils such as thymol, carvacrol, *p*-cymene and borneol (Gormez et al., 2016; Goze et al. 2009).

The isolation of 1-tetradecanol, rosmarinic acid and lithospermic acid from *O. rotundifolium* was recorded for the first time in this study.

**Esen Sezen Karaoglan<sup>1</sup>, Ufuk Özgen<sup>2</sup> and Cavit Kazaz<sup>3</sup>**

<sup>1</sup>Atatürk University, Faculty of Pharmacy, Department of Pharmaceutical Botany, 25240 Erzurum, Turkey; <sup>2</sup>Karadeniz Technical University,

Faculty of Pharmacy, Department of Pharmacognosy, 61080 Trabzon, Turkey; <sup>3</sup>Atatürk University, Faculty of Science, Department of Chemistry, 25240 Erzurum, Turkey.

Corresponding author:

email: esen.karaoglan@atauni.edu.tr

## References

- Aliγιannis N, Kalpoutzakis E, Mitaku S, Chinou IB. Composition and antimicrobial activity of the essential oils of two *Origanum* species. J Agric Food Chem. 2001; 49: 4168-70.
- Baser K, Ozek T, Tumen G, Sezik E. Composition of the essential oils of Turkish *Origanum* species with commercial importance. J Essent Oil Res. 1993; 5: 619-23.
- Baser KHC. The Turkish *Origanum* species. In: Oregano: The genera *Origanum* and *Lippia*. Kintzios SE (ed). London, Taylor and Francis, 1978, pp 109-26.
- Gormez A, Bozari S, Yanmis D, Gulluce M, Agar G, Sahin F. The use of essential oils of *Origanum rotundifolium* as antimicrobial agent against plant pathogenic bacteria. TEOP. 2016; 19: 656-63.
- Goze I, Alim A, Tepe AS, Sokmen M, Sevgi K, Tepe B. Screening of the anti-oxidant activity of essential oil and various extracts of *Origanum rotundifolium* Boiss. From Turkey. J Med Plants Res. 2009; 3: 246-54.
- Junges MJ, Fernandes JB, Vieira PC, Fernandes MDFGS, Filho ER, Frühauf M, Barañano AG. Triterpenosursânicos e oleanânicos isolados do caule de *Eugenia florida* DC. Revista de Pesquisa e Pós-Graduação. 2000; 1: 13-30.
- Kelley CJ, Harruff RC, Carmack M. Polyphenolics acids of *Lithospermum ruderale* II. carbon-13 nuclear magnetic resonance of lithospermic and rosmarinic acids. J Org Chem. 1976; 41: 449-55.
- Kelley CJ, Mahajan JR, Brooks LC, Neubert LA, Breneman WR, Carmack M. Polyphenolic acids of *Lithospermum ruderale* (Boraginaceae). I. Isolation and structure determination of lithospermic acid. J Org Chem. 1975; 40: 1804-15.
- Kotan R, Cakir A, Dadasoglu F, Aydin T, Cakmakci R, Ozer H, Kordali S, Mete E, Dikbas N. Antibacterial activities of essential oils and extracts of Turkish *Achillea*, *Satureja* and *Thymus* species against plant pathogenic bacteria. J Sci Food Agric. 2010; 90: 145-60.
- Novak J, Bitsch C, Langbehn J, Pank F, Skoula M, Gotsiou Y, Franz CM. Ratios of *cis*- and *trans*-sabinenehydrate in *Origanum majorana* L. and *Origanum microphyllum* (Benth.) Vogel. Biochem Syst Ecol. 2000; 28: 697-704.