


# Bangladesh Journal of Pharmacology

Volume: 12; Number 1; Year 2017



Cite this article as: Mohan K, Padmanaban AM, Uthayakumar V, Chandrasekhar R, Muralisankar T, Chithra E. Anti-cancer effect of the polysaccharide extract from the *Ganoderma lucidum* against HeLa cell lines. Bangladesh J Pharmacol. 2017; 12: 56-57.



## Letter to the Editor

### Anti-cancer effect of the polysaccharide extract from the *Ganoderma lucidum* against HeLa cell lines

Sir,

*Ganoderma lucidum* is an important Chinese medicinal mushroom which containing considerable amount of polysaccharides which has been reported to be effective as immunomodulation, antiviral, anti-oxidation, antibacterial, anti-cancer effects, etc., without toxic effects in animals (Zhang et al., 2002). It is known to possess various chemical compounds such as polysaccharides, triterpenoids, flavanoids, coumarins, quinones, carotenoids and amino acids as having antibacterial properties (Roberts, 2004). Polysaccharides are the best known and most potent mushroom-derived substances with antitumor and immunomodulating properties. Some recent studies described polysaccharide extracts of mushrooms, as *in vitro* inhibitors of various cancer cell lines such as colon, promyelocytic leukemia and gastric carcinoma (Masuda et al., 2009; Lavi et al., 2006;

Wong et al., 2007). The present study was found to investigate the anti-cancer effect of the polysaccharide extract from the *G. lucidum* against HeLa cell lines.

Matured *G. lucidum* fruit bodies were collected from the Ooty Hill region and authenticated by the Mycology Division of the Indian Forest Genetics and Tree Breeding Institute, Coimbatore, Tamilnadu, India. Fruiting bodies of mushrooms were dried at 45–50°C for 48 hours and powdered. The *G. lucidum* polysaccharides were isolated according to Chen et al. (2009). Fruiting bodies of mushrooms were dried at 45–50°C for 48 hours and powdered. The powdered material (200 g) was extracted with petroleum ether in Soxhlet apparatus for 8–10 hours. The extraction was done in four batches of 500 g each. The precipitate was collected by centrifugation (20 min at 9,000 g) and redissolved in double distilled water, and treated with Sevag's reagent several times to remove protein and then dialyzed against deionised water for 48 hours at 4°C. It was then evaporated at low temperature and lyophilized to obtain *G. lucidum* polysaccharide (900 mg) as a light brown powder. HeLa cell lines were purchased from

the National Centre for Cell Science, Pune, India. The anti-cancer effect of the polysaccharide extract of the *G. lucidum* was investigating using the MTT assay (Lau et al, 2004). The OD value was measured at 570 nm. The IC<sub>50</sub> value as the concentration of sample which reduced absorbance by 50% relative to the vehicle-treated control.

The proliferation of HeLa cell was significantly inhibited by *G. lucidum* polysaccharide. The Figure 1 and Table I shows the changes of the percentage of cell viability treated with *G. lucidum* polysaccharide extract (12.5, 25, 50, 100, and 200 µg/mL) in HeLa cell. There was 99% cell death at 200 µg/mL concentration was observed. The inhibi-




Figure 1: Cytomorphological alterations of HeLa cells treated with *G. lucidum* polysaccharide extract. A) Control (without treatment) indicated accumulation of HeLa cells uniformly. B, C, D, E, F) HeLa cells treated with 12.5, 25, 50, 100, 200 µg/mL polysaccharide extract for 48 hours indicate apoptotic features such as cytoplasmic blebbing (white arrows), round shape (orange arrows) and apoptotic body formation (black arrows), respectively. Magnification = x 400



Table I	
Effect of <i>G. lucidum</i> polysaccharide on HeLa cell lines	
Extract (µg/mL)	% Cell inhibition
12.5	2.5
25	5.0
50	7.4
100	24.1
200	99.4

tory concentration 50% (IC<sub>50</sub>) was fixed as 7.35 µg/mL. It shows a series of changes including cell shrinkage, cytoplasmic blebbing, and apoptotic body formation. *G. lucidum* polysaccharide that showed significant antitumor efficacy *in vitro* against HepG2, HeLa and A549 cancer cell lines, through both direct cytotoxic effects on tumor cells and growth-promoting effects on spleen cells (Li et al., 2010). From the present findings, it can be concluded that the *G. lucidum* polysaccharide extract shows toxicity to the HeLa cells.

**K. Mohan<sup>1</sup>, A. M. Padmanaban<sup>1</sup>, V. Uthayakumar<sup>1</sup>, R. Chandrasekhar<sup>1</sup>, T. Muralisankar<sup>2</sup> and E. Chithra<sup>1</sup>**

<sup>1</sup>Postgraduate and Research Department of Zoology, Sri Vasavi College, Erode, India; <sup>2</sup>Department of Marine Science, Bharathidasan University, Tiruchirappalli, India.

Corresponding author:  
email: kmohanphd@gmail.com

## References

Chen XP, Chen Y, Li SB, Chen YG, Lan JY, Liu LP. Free radical

scavenging of *Ganoderma lucidum* polysaccharides and its effect on antioxidant enzymes and immunity activities in cervical carcinoma rats. Carbohydr Polym. 2009; 77: 389-93.

Lau CBS, Ho CY, Kim CF, Leung KN, Fung KP, Tse TF. Cytotoxic activities of *Coriolus versicolor* (Yunzhi) extract on human leukemia and lymphoma cells by induction of apoptosis. Life Sci. 2004; 75: 797-808.

Lavi I, Friesem D, Geresh S, Hadar Y, Schwartz B. An aqueous polysaccharide extract from the edible mushroom *Pleurotus ostreatus* induces anti-proliferative and pro-apoptotic effects on HT-29 colon cancer cells. Cancer Lett. 2006; 244: 61-70.

Li N, Hu YL, He CX, Hu CJ, Zhou J, Tang GP, Gao JQ. Preparation, characterisation and anti-tumour activity of *Ganoderma lucidum* polysaccharide nanoparticles. J Pharm Pharmacol. 2010; 62: 139-44.

Masuda Y, Inoue M, Miyata A, Mizuno S, Nanba H. Maitake β-glucan enhances therapeutic effect and reduces myelosuppression and nephrotoxicity of cisplatin in mice. Int Immunopharmacol. 2009; 9: 620-26.

Roberts LM. Australian Ganoderma: Identification, growth and antibacterial properties. PhD. Thesis, Swinburne University of Technology, Australia, 2004.

Wong SM, Wong KK, Chiu LCM, Cheung PCK. Non-starch polysaccharides from different developmental stages of *Pleurotus tuber-reginum* inhibited the growth of human acute promyelocytic leukemia HL-60 cells by cell-cycle arrest and/or apoptotic induction. Carbohydr Polym. 2007; 68: 206-17.

Zhang GL, Wang YH, Ni W, Teng HL, Lin Z. Hepatoprotective role of *Ganoderma lucidum* polysaccharide against BCG-induced immune liver injury in mice. World J Gastroenterol. 2002; 8: 728-33.