



**BJP**

**Bangladesh Journal of Pharmacology**

**Research Article**

**Isolation and antifungal screening  
of endophytic fungi from *Erigeron  
canadensis***

## Isolation and antifungal screening of endophytic fungi from *Erigeron canadensis*

Xuelian Bai<sup>1</sup>, Ting Zhou<sup>1</sup>, Tongfei Lai<sup>1</sup>, Yicong Li<sup>1</sup>, Jiale Chai<sup>1</sup>, Jiajun Ni<sup>1</sup> and Huawei Zhang<sup>2</sup>

<sup>1</sup>College of Life and Environmental Sciences, Hangzhou Normal University, Hangzhou 310036, P. R. China;

<sup>2</sup>School of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, P. R. China.

### Article Info

Received: 9 April 2017  
Accepted: 5 June 2017  
Available Online: 16 July 2017  
DOI: 10.3329/bjp.v12i3.32126

#### Cite this article:

Bai X, Zhou T, Lai T, Li Y, Chai J, Ni J, Zhang H. Isolation and antifungal screening of endophytic fungi from *Erigeron canadensis*. Bangladesh J Pharmacol. 2017; 12: 256-59.

### Abstract

Sixteen fungal strains isolated from the *Erigeron canadensis*, one of traditional Chinese medicines used to treat the pathogenic infection and dysentery, were evaluated for their antifungal activities against one human pathogen *Candida albicans*, and two phytopathogens, *Colletotrichum fructicola* and *Rhizoctonia cerealis*. The bioassay results indicated that the ethyl acetate extract of the fermentation broth of these fungal endophytes had stronger antimicrobial activities. Among these endophytic strains, the ethyl acetate extracts of strains NPR003 and NPR005 showed the strongest inhibitory effects and has potential application in the discovery of new antifungal agents. This was the first report on the isolation of endophytic fungi from *E. canadensis* and evaluation of their antifungal activities.

## Introduction

Fungal infection is an important life-threatening disease, which annually causes the death of millions of people (Xie et al., 2014) and the great loss of crop production (Bennett et al., 2012). Effective and efficient control of fungal pathogens can be achieved by the use of chemically synthetic fungicides. However, a limited repertoire of eco-friendly and low toxic antifungals are used all over the world. Thus, there is an urgent need to search for alternative approaches to accelerate the discovery of more biocontrol agents to treat fungal infections and their drug resistance. A growing evidence indicates that endophyte, a special microbial community colonized in healthy plants, is one of the rich sources of bioactive natural products with great potential application in the fields of medicine and agriculture (Zhang et al., 2006; 2014). Isolation and antifungal evaluation of endophytic microbe would provide a better alternative way to identify novel antifungal agents.

*Erigeron canadensis* Linn., called horseweed widely distributed in China, is one of the herbaceous Chinese medicines used to treat intestinal diseases (Park et al., 2013; Wu et al., 2015). Modern pharmacological study showed that *E. canadensis* has strong bioactivities, including anti-inflammation (Sohn et al., 2009; Sung et al., 2014), anticoagulant and antiplatelet activity (Pawlacyk et al., 2011), inhibition of melanogenesis (Hong et al., 2008), cytotoxicity (Choi et al., 2008) and antifungal activity (Curini et al., 2003). Up to now, however, there is no report on biology and chemistry of endophytic microbes from this medicinal plant. In our continuous investigation of antimicrobial fungi derived from the plants (Zhang et al., 2012; 2014; 2015), 16 endophytic strains (numbered as NPR001-NPR016) were isolated from *E. canadensis* and evaluation of their antifungal effects was carried out in this work.

## Materials and Methods

The healthy plant of *E. canadensis* was obtained from the



Xiasha campus of Hangzhou Normal University, Hangzhou, China and used for the endophyte isolation within 48 hours after harvest. Three testing fungal pathogens, including one human pathogen, *Candida albicans*, and two plant pathogens, *Colletotrichum fructicola* and *Rhizoctonia cerealis*, were from China Center for Type Culture Collection (CCTCC). All chemicals used in this project were of analytical grade.

#### Isolation of endophytic strain

Endophytic fungal strains from *E. canadensis* were isolated according to our reported procedure (Zhang et al., 2012; 2014; 2015). All these fungal strains were transferred into potato dextrose agar (PDA) slants followed by storing at 4°C.

#### Fermentation and preparation of ethyl acetate extract of fungal endophyte

Each fungal strain was cultured on PDA at 28°C for 7 days. Then a balanced amount of fungal colony was transferred to culture broth in a 500 mL erlenmeyer flask, which contained 200 mL sterilized potato dextrose broth (PDB) followed by shaking at 150 rpm for 15 days at 28°C. By the end of fermentation, the mycelium and the culture broth were separated by centrifugation at 4,500 rpm for 15 min at 4°C. The fermentation broth was extracted twice with 400 mL ethyl acetate (Merck). Then the upper solvent was evaporated at 20°C in vacuum to yield the ethyl acetate extract, which possibly had antimicrobial substances metabolized by fungal endophyte. Each afforded extract was kept in a vacuum drier for 3 days and dissolved in dimethyl sulfoxide (DMSO, Merck) before bioassay. The final concentration of each ethyl acetate extract had three levels, including 0.1, 1, 10 mg/mL.

#### Antifungal test

The antifungal assay was carried out using disc diffusion method (Zhang et al., 2012; 2014; 2015). Three fungal blocks of *C. albicans*, *C. fructicola* and *R. cerealis* were respectively transferred into three 500 mL erlenmeyer flasks, each flask had 200 mL sterilized PDB. The seed liquid was prepared after incubation at 28°C on a rotary shaker at 150 rpm for 3 days. Firstly, 5 mL melt water agar (WA) medium was evenly poured into petri dishes ( $\Phi = 90$  mm). Secondary, 200  $\mu$ L seed liquid was added to fresh NA medium and mixed well. Thirdly, the same amount of fresh NA medium was poured on the solidified WA medium and the testing plate with double medium for antifungal bioassay was prepared. After 5 holes ( $\Phi = 6$  mm) were equidistantly drilled on inoculated media, a piece of standard sterile filter paper ( $\Phi = 6$  mm) was put in one hole followed by adding 100  $\mu$ L ethyl acetate extract of endophytic fungal strain. Amphotericin B (30  $\mu$ g/disk, Sigma-Aldrich) was used as the positive control and the pure DMSO or sterilized water was the negative control. The

diameter of inhibition zone (in mm) was measured to assess antifungal activity. All tests were carried out in triplicate.

---

## Results

Biological investigation of endophytic fungi from healthy *E. canadensis* lead to isolation of sixteen fungal strains (NPR001-NPR016), which seven strains NPR001-NPR007 were from roots, two strains NPR008-NPR009 from flowers, five strains NPR010-NPR014 from stems, and two strains NPR015-NPR016 from leaves. The antifungal bioassay results were shown in Table I.

The ethyl acetate extracts of endophytic fungi had different inhibitory effect on fungal pathogens, including *C. albicans*, *C. fructicola* and *R. cerealis*. Generally, their antifungal activities were in a concentration-dependent manner. Among these fungal endophytes, two strains NPR003 and NPR005 had a broad spectrum of antifungal activities against all testing pathogens. As far as the human fungal pathogen *C. albicans* concerned, 7 strains NPR003-NPR005, NPR007, NPR009, NPR012 and NPR013 exhibited stronger inhibitory effect than others at the same concentration. Five endophytic strains (NPR003, NPR005, NPR006, NPR010 and NPR015) and 2 strains (NPR001 and NPR005) showed potent activity against *C. fructicola* and *R. cerealis*, respectively.

---

## Discussion

According to symbiosis theory, endophytic microbe has the ability to biosynthesize some 'phytochemicals' originally from their host plants (Zhang et al., 2014). A growing evidence suggests that fungal endophytes are widely distributed in healthy plant organs and have abundant biodiversity and chemodiversity. Bioactive extracts and natural products derived from endophytic fungi would effectively accelerate new drug discovery and contribute to the development of green agriculture.

In the present work, sixteen fungal strains (coded as NPR001-NPR016) were characterized from *E. canadensis* Linn. Bioassay results showed that the ethyl acetate extracts of strains NPR003 and NPR005 exhibited potent inhibitory effects on all test pathogens, which were in a concentration-dependent manner. It also indicated that strains NPR003 and NPR005 were the best biocontrol candidates and had potential application in medicine and pesticide industry.

Up to now, strains NPR003 and NPR005 were respectively identified as *Alternaria* spp., *Aspergillus* spp. by 18S rDNA sequence analysis and morphological properties. Bioassay-guided fractionation of the ethyl acetate extracts of the scale-up fermentation broth of

Table I				
Antifungal activities of ethyl acetate extracts of endophytic fungi from <i>Erigeron canadensis</i>				
Strain No.	Concentration mg/mL	Antifungal effects		
		<i>Candida albicans</i>	<i>Colletotrichum fructicola</i>	<i>Rhizoctonia cerealis</i>
NPR001	0.1	-	-	-
	1	-	-	+
	10	+	+	++
NPR002	0.1	-	-	-
	1	+	-	-
	10	+	+	+
NPR003	0.1	+	+	-
	1	++	++	+
	10	+++	++	++
NPR004	0.1	-	-	-
	1	+	-	+
	10	++	-	+
NPR005	0.1	+	+	+
	1	++	++	++
	10	+++	+++	+++
NPR006	0.1	-	-	-
	1	-	+	-
	10	-	++	+
NPR007	0.1	+	-	-
	1	++	-	-
	10	++	+	-
NPR008	0.1	-	-	-
	1	-	-	-
	10	+	-	+
NPR009	0.1	+	-	-
	1	++	-	-
	10	++	+	+
NPR010	0.1	-	-	-
	1	-	+	+
	10	-	++	+
NPR011	0.1	-	-	-
	1	-	-	-
	10	+	+	-
NPR012	0.1	-	-	-
	1	++	+	-
	10	++	+	+
NPR013	0.1	+	-	-
	1	+	-	+
	10	++	+	+
NPR014	0.1	-	-	-
	1	-	-	-
	10	-	-	-
NPR015	0.1	+	-	-
	1	+	+	-
	10	+	++	+
NPR016	0.1	-	-	-
	1	-	-	-
	10	+	-	+
Amphotericin B	30 µg/disk	++++	++++	++++
DMSO		-	-	-

<sup>a</sup>Expressed by the diameter of inhibition zones: -, no inhibition; +, <10 mm; ++, 10-15 mm; +++, 16-20 mm; +++++, >20 mm



strain NPR003 led to the isolation of five tricycloalternarene (TCA) analogs, which had anti-microbial and cytotoxic effects (Wang et al., 2013). Our findings would supply an important reference to discover a new source of TCA compounds as antimicrobial agents.

## Conclusion

This was the first report on isolation and evaluation of antimicrobial effect of endophytic fungi from *E. canadensis*.

## Financial Support

Innovation and Entrepreneurship Program for National Student (201610346024), the Tri-Five Cultivating Project (2016XJSGWXM26)

## Conflict of Interest

Authors have declare no conflict of interest

## Acknowledgements

The Competition Project for 17<sup>th</sup> National Student Challenge Cup of Extracurricular Academic Science and Technology Works from Hangzhou Normal University were gratefully appreciated.

## References

Bennett AJ, Bending GD, Chandler D, Hilton S, Mills P. Meeting the demand for crop production: The challenge of yield decline in crops grown in short rotation. *Biol Rev.* 2012; 87: 52-71.

Choi HJ, Wang HY, Kim YN, Heo SJ, Kim NK, Jeong MS, Park YH, Kim S. Composition and cytotoxicity of essential oil extracted by steam distillation from horseweed (*Erigeron canadensis* L.) in Korea. *J Korean Soc Appl Biol Chem.* 2008;

51: 55-59.

- Curini M, Bianchi A, Epifano F, Bruni R, Torta L, Zambonelli A. Composition and *in vitro* antifungal activity of essential oils of *Erigeron canadensis* and *Myrtus communis* from France. *Chem Nat Compd.* 2003; 39: 191-94.
- Hong ES, Nguyen DTM, Nguyen DH, Kim EK. Inhibition of melanogenesis by *Erigeron canadensis* via down-regulating melanogenic enzymes in B16F10 melanoma cells. *Korean J Chem Eng.* 2008; 25: 1463-66.
- Park WS, Bae JY, Chun MS, Chung HJ, Han SY, Ahn MJ. Suppression of gastric ulcer in mice by administration of *Erigeron canadensis* extract. *Proc Nutr Soc.* 2013; 72: E263-63.
- Pawlaczyk I, Czerchawski L, Kuliczkowski W, Karolko B, Pilechi W, Witkiewicz W, Gancarz R. Anticoagulant and anti-platelet activity of polyphenolic-polysaccharide preparation isolated from the medicinal plant *Erigeron canadensis* L. *Thromb Res.* 2011; 127: 328-40.
- Sohn SH, Ko E, Oh BG, Kim J, Choi E, Kim SH, Kim Y, Shin M, Hong M, Bae H. Global gene analysis of *Erigeron canadensis*-treated TNF-alpha-, IL-4- and IL-1 beta-stimulated A549 human epithelial cells. *Ann Nutr Metab.* 2009; 54: 227-35.
- Sung J, Sung M, Kim Y, Ham H, Jeong HS, Lee J. Anti-inflammatory effect of methanol extract from *Erigeron canadensis* L. may be involved with up-regulation of heme oxygenase-1 expression and suppression of NF kappa B and MAPKs activation in macrophages. *Nutr Res Pract.* 2014; 8: 352-59.
- Wu CY, Shao S, Yan MM, Fu ML, Xu DM. Research on development chemical constituents of *Erigeron canadensis* L. *China Pharm.* 2015; 24: 1-3.
- Xie JL, Polvi EJ, Shekhar-Guturja T, Cowen LE. Elucidating drug resistance in human fungal pathogens. *Future Microbiol.* 2014; 9: 523-42.
- Zhang HW, Bai XL, Wu BX. Evaluation of antimicrobial activities of extracts of endophytic fungi from *Artemisia annua* Linn. *Bangladesh J Phamacol.* 2012; 7: 249-57.
- Zhang HW, Ruan CF, Bai XL. Isolation and antimicrobial effects of endophytic fungi from *Edgeworthia chrysantha*. *Bangladesh J Pharmacol.* 2015; 10: 529-32.

### Author Info

Huawei Zhang (Principal contact)  
e-mail: hwzhang@zjut.edu.cn