

# Mydriatic eye drop

(Pre- and post-test)

1. Which of the following you should not do before instilling eye drop?
  - a) Checking the name of medication
  - b) Looking for the expire date
  - c) Writing down the date of opening in the label
  - d) Touching the tip of eye drop bottle for finding any slit
2. In case of selection of a mydriatic eye drop to perform fundoscopic examination of eye, which of the following point you need not consider?
  - a) Efficacy and safety
  - b) Cost-effectiveness
  - c) Color of the cap of the eye drop
  - d) Brand name
3. What should you do after instill a drop of medication in eye?
  - a) Keep looking upward without blinking
  - b) Close your eyes firmly
  - c) Give a gentle pressure with index finger at the junction of nasal bridge & inner side of eye
  - d) Rinse eye with water
4. What should be the gap to instill a second drop in same eye?
  - a) 1 minute
  - b) 3 minutes
  - c) 5 minutes
  - d) 7 minutes
5. Which method should be followed to instill eye drop in a person with poor vision?
  - a) Tactile
  - b) Taking help of other
  - c) None of the above
  - d) Both 'a' and 'b'

6. Which of the following can make eye drop instillation easier?
  - a) Eye drop holder
  - b) Tactile method
  - c) Taking help of another person
  - d) Using mirror
  
7. After opening how long an eye drop bottle can be used?
  - a) 10 days
  - b) 14 days
  - c) 20 days
  - d) 28 days
  
8. Which of the following is the bottle top color of a mydriatic eye drop bottle?
  - a) Yellow
  - b) Orange
  - c) Red
  - d) Purple
  
9. In tactile method index finger is used as a guide to instill an eye drop. T/F
  
10. After instilling a drop of medication in eye, give a gentle pressure at junction of nasal bridge and inner side of eye for 1-2 minutes. T/F